arXiv:math/0703585v1 [math.CO] 20 Mar 2007

Maxmaxflow and Counting Subgraphs

Bill Jackson
School of Mathematical Sciences
Queen Mary, University of London
Mile End Road
London E1 4NS, England
`B.JACKSON@QMUL.AC.UK`

Alan D. Sokal*
Department of Physics
New York University
4 Washington Place
New York, NY 10003 USA
`SOKAL@NYU.EDU`

15 March 2007

Abstract

We introduce a new graph invariant $\Lambda(G)$ that we call maxmaxflow, and put it in the context of some other well-known graph invariants, notably maximum degree and its relatives. We prove the equivalence of two "dual" definitions of maxmaxflow: one in terms of flows, the other in terms of cocycle bases. We then show how to bound the total number (or more generally, total weight) of various classes of subgraphs of G in terms of either maximum degree or maxmaxflow. We conclude with some conjectures concerning the relevance of maxmaxflow to the roots of chromatic polynomials.

Key Words: Graph, subgraph, flow, cocycle, maxmaxflow, maximum degree, second-largest degree, degeneracy number, chromatic polynomial.

Mathematics Subject Classification (MSC) codes: 05C99 (Primary); 05C15, 05C30, 05C35, 05C40, 82B20, 90B10 (Secondary).

---

*Also at Department of Mathematics, University College London, London WC1E 6BT, England.

Contents

1 Introduction **3**

2 Cocycle Bases and Maxmaxflow **7**

3 Some Preliminaries **11**
3.1 Pointwise bounds vs. generating-function bounds 12
3.2 Convex hulls in graphs 12

4 Counting Walks and Paths **13**

5 Counting Trees and Forests **19**
5.1 The classes $\mathcal{T}_m(X)$ and $\mathcal{F}_m(X,Y)$ 19
5.2 The class $\mathcal{H}_m(X)$ 23

6 Counting Connected Subgraphs **24**

7 Counting Blocks, Block Paths, Block Trees and Block Forests **30**
7.1 The classes $\mathcal{BT}_m(X)$, $\mathcal{BF}_m(X,Y)$ and $\mathcal{BF}^*_m(X,Y)$ 30
7.2 The class $\mathcal{B}_m(X)$ 42

8 A Conjecture on Chromatic Roots **45**

1 Introduction

The purpose of this paper is to introduce a new graph invariant $\Lambda(G)$ that we call *maxmaxflow*, to put it in the context of some other well-known graph invariants (notably maximum degree and its relatives), and to use it to prove upper bounds on the numbers of subgraphs of certain types.

Let G be a finite undirected graph with vertex set $V(G)$ and edge set $E(G)$; in this paper all graphs are assumed to be loopless, but multiple edges are allowed unless explicitly specified otherwise. We shall say that G is *simple* if it has no multiple edges. Let $\Delta(G) = \max_{x \in V(G)} d_G(x)$ be the maximum degree of G, and more generally let $\Delta_k(G)$ be the kth largest degree of G:

$$\Delta_k(G) \;=\; \min_{x_1,\ldots,x_{k-1} \in V(G)} \quad \max_{x \in V(G) \setminus \{x_1,\ldots,x_{k-1}\}} d_G(x) \;. \tag{1.1}$$

We trivially have

$$\delta(G) \;\equiv\; \Delta_n(G) \;\le\; \cdots \;\le\; \Delta_3(G) \;\le\; \Delta_2(G) \;\le\; \Delta_1(G) \;\equiv\; \Delta(G) \tag{1.2}$$

where $n = |V(G)|$. A special role will be played in this paper by the second-largest degree, $\Delta_2(G)$.

For $x, y \in V(G)$ with $x \neq y$, *the maximum flow* from x to y in G is

$$\begin{aligned} \lambda_G(x,y) \;&=\; \max \text{ \# of edge-disjoint paths from } x \text{ to } y && \text{(1.3a)} \\ &=\; \min \text{ \# of edges separating } x \text{ from } y && \text{(1.3b)} \end{aligned}$$

We then define the *maxmaxflow* of G

$$\Lambda(G) \;=\; \max_{\substack{x,\, y \,\in\, V(G) \\ x \neq y}} \lambda_G(x,y) \;. \tag{1.4}$$

[Note the contrast with the edge-connectivity, which is the *minimum* of $\lambda_G(x,y)$ over $x \neq y$.] Clearly $\lambda_G(x,y) \le \min[d_G(x), d_G(y)]$, so that

$$\Lambda(G) \;\le\; \Delta_2(G) \;. \tag{1.5}$$

We will show later (Proposition 2.10) that $\Lambda(G) \ge \Delta_{n-1}(G)$. Note that several cases can arise:

(a) $\Lambda(G) = \Delta_2(G) = \Delta(G)$. Indeed, in any regular graph one has $\Lambda(G) = \Delta_i(G)$ for all i $(1 \le i \le n)$.

(b) $\Lambda(G) = \Delta_2(G) \ll \Delta(G)$. This occurs, for example, in stars $K_{1,r}$ and wheels $K_1 + C_r$.

(c) More generally, one can have $\Lambda(G) = \Delta_{j+1}(G) \ll \Delta_j(G)$ for any fixed integer j. Moreover, such examples can be taken to be k-connected for arbitarily large k.[1]

Note also that maxmaxflow has a naturalness property that maximum degree and kth-largest degree lack, namely, it "trivializes over blocks": $\Lambda(G) = \max_{1 \le i \le b} \Lambda(G_i)$ where $G_1, \ldots, G_b$ are the blocks of G (Proposition 2.11).

Maxmaxflow appears to have been considered sporadically in the graph-theoretic literature. Bollobás [2, section I.5] addresses some extremal problems involving maxmaxflow in simple graphs (he uses the term "maximum local edge-connectivity" and denotes it $\bar{\lambda}(G)$); see likewise Mader [9, section IV]. In particular, Mader [8] has shown that whenever an n-vertex graph has more than $k(n-1)/2$ edges, it has maxmaxflow at least k, but that for every $n \ge k \ge 2$ there exists an n-vertex graph with exactly $\lfloor k(n-1)/2 \rfloor$ edges and maxmaxflow $k-1$.[2]

An apparently very different quantity can be defined via cocycle bases. For X, Y disjoint subsets of $V(G)$, let $E(X,Y)$ denote the set of edges in G between X and Y. A *cocycle* of G is a set $E(X, X^c)$ where $X \subseteq V(G)$ and $X^c \equiv V(G) \backslash X$. It is well-known that the cocycles of G form a vector space over GF(2) with respect to symmetric difference; this is called the *cocycle space* of G. Let $\widetilde{\Lambda}(G)$ be the minmax cardinality of the cocycles in a basis, i.e.

$$\widetilde{\Lambda}(G) \;=\; \min_{\mathcal{B}} \max_{C \in \mathcal{B}} |C| \tag{1.6}$$

where the min runs over all bases $\mathcal{B}$ of the cocycle space of G. Since one special class of cocycle bases consists of taking the stars $C(x) = E(\{x\}, \{x\}^c)$ for all but one of the vertices in each component of G, we clearly have

$$\widetilde{\Lambda}(G) \;\le\; \Delta_2(G)\;. \tag{1.7}$$

The relationship, if any, between maxmaxflow and cocycle bases is perhaps not obvious at first sight. But we shall prove (Corollary 2.9) that

$$\Lambda(G) \;=\; \widetilde{\Lambda}(G)\;. \tag{1.8}$$

---

[1] Proof. For $1 \le i \le j$, let H_i be a k-connected graph with one vertex v_i of degree $\Delta \gg k$ and all other vertices of degree k. [Such graphs can be constructed by taking a $(k-1)$-connected $(k-1)$-regular graph with a large number Δ of vertices and adding a new vertex v_i adjacent to every other vertex.] Construct G from the disjoint union of $H_1, H_2, \ldots, H_j$ by adding k edges between each pair $H_i - v_i$ and $H_{i+1} - v_{i+1}$ $(1 \le i \le j-1)$ in such a way that the set of edges of G which do not belong to any H_i are independent. [This can be done as long as $|V(H_i)| \ge 2k+1$.] Then G is k-connected and satisfies $\Lambda(G) = \Delta_{j+1}(G) = k+1$. [Since all pairs of vertices of G with degrees greater than $k+1$ are of the form v_s, v_t with $s \ne t$, and hence are separated by a set of k edges, we have $\Lambda(G) \le k+1$. On the other hand, if we choose two vertices $x, y \in H_1$ that are both adjacent to H_2, we can find k edge-disjoint xy-paths in H_1 (since H_1 is k-connected) and an extra xy-path passing through H_2.] But $\Delta_j(G) = \Delta$.

[2] For $k = 2, 3$ this is easy. For $k = 4$ it was proven earlier by Bollobás [1], and for $k = 5, 6$ by Leonard [5, 6].

The two definitions thus give dual approaches to the same quantity.

Finally, define the *degeneracy number* $D(G) = \max_{H \subseteq G} \delta(H)$, where the max runs over all subgraphs H of G, and $\delta(H)$ is the minimum degree of H. It is easy to see that

$$D(G) \;\le\; \Delta_2(G) \tag{1.9}$$

[if H has at least two vertices, then $\delta(H) \le \Delta_2(G)$; otherwise $\delta(H) = 0$]. But we shall in fact show (Proposition 2.10) that

$$D(G) \;\le\; \Lambda(G) \;. \tag{1.10}$$

In summary, therefore, we have

$$D(G) \;\le\; \Lambda(G) \;=\; \widetilde{\Lambda}(G) \;\le\; \Delta_2(G) \;\le\; \Delta(G) \;. \tag{1.11}$$

The natural setting for the results of this paper is, in fact, that of a finite undirected loopless (multi)graph G equipped with nonnegative real edge weights $\mathbf{w} = \{w_e\}_{e \in E(G)}$. Indeed, all of the aforementioned invariants have natural generalizations to this context. Define first the weighted degree of a vertex,

$$d_G(x, \mathbf{w}) \;=\; \sum_{e \ni x} w_e \;. \tag{1.12}$$

We then set

$$\begin{aligned}
\Delta(G, \mathbf{w}) &= \max_{x \in V(G)} d_G(x, \mathbf{w}) && (1.13) \\
\Delta_k(G, \mathbf{w}) &= \min_{x_1, \ldots, x_{k-1} \in V(G)} \;\; \max_{x \in V(G) \setminus \{x_1, \ldots, x_{k-1}\}} d_G(x, \mathbf{w}) && (1.14) \\
\delta(G, \mathbf{w}) &= \min_{x \in V(G)} d_G(x, \mathbf{w}) && (1.15) \\
\widetilde{\Lambda}(G, \mathbf{w}) &= \min_{\mathcal{B}} \max_{C \in \mathcal{B}} \sum_{e \in C} w_e && (1.16) \\
D(G, \mathbf{w}) &= \max_{H \subseteq G} \delta(H, \mathbf{w}|_H) && (1.17)
\end{aligned}$$

Likewise, max-flow quantities are naturally defined when the $\{w_e\}$ are interpreted as edge capacities:

$$\begin{aligned}
\lambda_G(x, y; \mathbf{w}) &= \text{max flow from } x \text{ to } y \text{ with edge capacities } \mathbf{w} && (1.18a) \\
&= \text{min cut between } x \text{ and } y \text{ with edge capacities } \mathbf{w} && (1.18b)
\end{aligned}$$

and thence

$$\Lambda(G, \mathbf{w}) \;=\; \max_{\substack{x, y \in V(G) \\ x \neq y}} \lambda_G(x, y; \mathbf{w}) \;. \tag{1.19}$$

In this generality we shall prove

$$D(G, \mathbf{w}) \;\le\; \Lambda(G, \mathbf{w}) \;=\; \widetilde{\Lambda}(G, \mathbf{w}) \;\le\; \Delta_2(G, \mathbf{w}) \;\le\; \Delta(G, \mathbf{w})\,. \tag{1.20}$$

The unweighted case corresponds to setting all edge weights to 1.

Let us make a remark about the treatment of multiple edges. It is easy to see that all the quantities appearing in (1.20) are unchanged if we replace a family $e_1, \ldots, e_n$ of parallel edges with weights $w_{e_1}, \ldots, w_{e_n}$ by a single edge e with weight $w_e = \sum_{i=1}^{n} w_{e_i}$. So, in proving (1.20), we could, if we wanted, restrict attention to simple graphs; but we don't bother, because no simplification of the proof is obtained by doing so. Likewise, the weighted counts discussed in Sections 4 and 5 are unchanged by this replacement, because the subgraphs in question (walks, paths, trees and forests) can include at most one edge from a family of parallel edges. So it would suffice to prove the bounds in Sections 4 and 5 for simple graphs; but once again, we refrain from making this assumption because nothing is gained by doing so. For the weighted counts discussed in Sections 6 and 7, by contrast, no simple reduction of multiple edges can be performed, because the subgraphs in question *do* permit the inclusion of multiple edges. We shall therefore have to deal there with multigraphs in all our arguments.

Our interest in maxmaxflow was sparked initially by our study of the roots of chromatic polynomials [12]. It is an elementary fact [3, Theorem V.1] that all of the *integer* chromatic roots lie in the interval $[0, D(G)]$, i.e. the chromatic number $\chi(G)$ is at most $D(G) + 1$. On the other hand, Thomassen has shown that the *real* chromatic roots cannot be bounded in terms of the degeneracy number: indeed, there are 2-degenerate graphs with arbitrarily large real chromatic roots [14, Theorem 3.9] (see also [13, Appendix B]). Nevertheless, one of us has shown that *all* the chromatic roots (real or complex) can be bounded in terms of the second-largest degree: they lie in the disc $|q| < 7.963907\Delta_2(G) + 1$ [12, Corollary 6.4]. Furthermore, he conjectured, following a suggestion of Shrock and Tsai [10, 11], that it might be possible to bound all the chromatic roots in terms of maxmaxflow. An important step in [12] is obtaining a reasonably tight upper bound in terms of maximum degree for the number of connected m-edge subgraphs containing a fixed vertex. Our work here on counting subgraphs in terms of maxmaxflow is motivated in part by the goal of adapting the methods of [12] to this case.

The plan of this paper is as follows: In Section 2 we analyze cocycle bases and prove the fundamental result (1.20); an important role in this proof is played by *Gomory–Hu trees* [4]. The remainder of the paper is devoted to bounding the total number (or more generally, total weight) of various classes of subgraphs in terms of either maximum degree or maxmaxflow. Our basic approach is to start with a bound (sometimes a known one, sometimes a new one) in terms of maximum degree, and then see whether we can find a similar bound in terms of maxmaxflow. After some brief preliminaries (Section 3), we analyze walks and paths (Section 4) and then trees and forests (Section 5). In Section 6 we consider connected subgraphs and related objects, and in Section 7 we consider blocks, block paths, block trees and block forests. We close with a brief discussion of the conjectured relevance of maxmaxflow to the roots of chromatic polynomials (Section 8).

2 Cocycle Bases and Maxmaxflow

Given a graph G and disjoint subsets $X, Y \subseteq V(G)$, let $E(X, Y)$ denote the set of edges in G between X and Y. A *cocycle* of G is a set $E(X, Y)$ where X, Y is a bipartition of $V(G)$; note that $X = \varnothing$ and $Y = \varnothing$ are allowed. Let $\oplus$ denote symmetric difference. The following lemma is well known:

Lemma 2.1 *Let $C_1 = E(X_1, Y_1)$ and $C_2 = E(X_2, Y_2)$ be two cocycles in G. Then $C_1 \oplus C_2 = E\left((X_1 \cap X_2) \cup (Y_1 \cap Y_2),\, (X_1 \cap Y_2) \cup (Y_1 \cap X_2)\right)$.*

It follows that the set of all cocycles of G forms a vector space over GF(2) with respect to symmetric difference. This is the *cocycle space* of G. Its dimension is $|V(G)| - c(G)$, where $c(G)$ denotes the number of components of G.

Lemma 2.2 *Let G be a connected graph and let C be a cocycle of G. Then C corresponds to a unique bipartition of $V(G)$.*

Proof. Suppose $C = E(X_1, Y_1) = E(X_2, Y_2)$. Since $C \oplus C = \varnothing$ there are no edges in G from $(X_1 \cap X_2) \cup (Y_1 \cap Y_2)$ to $(X_1 \cap Y_2) \cup (Y_1 \cap X_2)$. Since G is connected, it follows that either $(X_1 \cap X_2) \cup (Y_1 \cap Y_2) = \varnothing$ and hence $(X_1, Y_1) = (Y_2, X_2)$, or else $(X_1 \cap Y_2) \cup (Y_1 \cap X_2) = \varnothing$ and hence $(X_1, Y_1) = (X_2, Y_2)$. $\square$

Lemma 2.3 *Let G be a connected graph, let C_1, C_2 be cocycles of G, and let x, y be vertices of G. Suppose that x, y belong to the same subset in the bipartitions of G corresponding to C_1 and C_2, respectively. Then x, y belong to the same subset in the bipartition of G corresponding to $C_1 \oplus C_2$.*

Proof. Immediate from Lemma 2.1. $\square$

Lemma 2.4 *Let G be a connected graph and let $C_1, C_2, \ldots, C_m$ be cocycles of G. Suppose that for each i, there exists a pair of vertices x_i, y_i such that x_i, y_i belong to different subsets in the bipartition of G corresponding to C_i and to the same subset in the bipartition of G corresponding to C_j for all $j \neq i$ $(1 \leq j \leq m)$. Then $C_1, C_2, \ldots, C_m$ are linearly independent.*

PROOF. If not, then we may suppose without loss of generality that $C_1 = C_2 \oplus C_3 \oplus \ldots \oplus C_m$. This contradicts the fact that x_1, y_1 belong to the different subsets in the bipartition corresponding to C_1 and to the same subset in the bipartition corresponding to $C_2 \oplus C_3 \oplus \ldots \oplus C_m$, by Lemma 2.3. $\square$

Let G be a connected graph and let T be a tree on the same vertex set V as G. (We emphasize that T need not be a subgraph of G.) Each edge $e \in E(T)$ induces a bipartition of V into nonempty subsets X, Y given by the two components of $T - e$; we define the *elementary cocycle of G corresponding to e and T* to be the cocycle $E_G(X, Y)$.

Lemma 2.5 *Let G be a connected graph with n vertices, and let T be a tree on the same set of vertices (not necessarily a subgraph of G). For each edge $e_i \in E(T)$, let C_i be the elementary cocycle of G corresponding to e_i and T. Then $\{C_1, C_2, \ldots, C_{n-1}\}$ is a basis for the cocycle space of G.*

PROOF. Using Lemma 2.4 (taking x_i, y_i to be the end-vertices of e_i) we deduce that $C_1, C_2, \ldots, C_{n-1}$ are linearly independent. Since the dimension of the cocycle space of G is $n - 1$, they form a basis. $\square$

Lemma 2.6 *Let G be a connected graph with n vertices, let $\{C_1, C_2, \ldots, C_{n-1}\}$ be a basis for the cocycle space of G, and let $x, y \in V(G)$ with $x \neq y$. Then x, y belong to different subsets in the bipartition corresponding to C_i, for some $1 \leq i \leq n - 1$.*

PROOF. Suppose not. Let C be a cocycle in G that separates x and y [for example, $E(\{x\}, \{x\}^c)$]. Since $\{C_1, C_2, \ldots, C_{n-1}\}$ is a basis for the cocycle space of G, C is a linear combination of $C_1, C_2, \ldots, C_{n-1}$. This contradicts Lemma 2.3. $\square$

Now let G be equipped with a family of nonnegative real edge weights $\mathbf{w} = \{w_e\}_{e \in E(G)}$. As in (1.18)/(1.19), we let $\lambda_G(x, y; \mathbf{w})$ be the max flow from x to y with edge capacities $\mathbf{w}$, and $\Lambda(G, \mathbf{w})$ the corresponding maxmaxflow. As in (1.16), we let $\widetilde{\Lambda}(G, \mathbf{w})$ be the minmax weight of the cocycles in a basis. In order to prove the fundamental result (1.20), we shall need the following classic result on flows (see [7, Section 2.3] for an excellent exposition):

Theorem 2.7 (Gomory and Hu [4]) *Let G be a connected graph equipped with nonnegative real edge weights $\mathbf{w} = \{w_e\}_{e \in E(G)}$. Then there exists a tree T with vertex set $V(T) = V(G) \equiv V$ (note that T is not necessarily a subgraph of G!) and a set $\mathbf{w}^T = \{w_e^T\}_{e \in E(T)}$ of nonnegative real edge weights such that*

(a) $\lambda_G(x, y; \mathbf{w}) = \lambda_T(x, y; \mathbf{w}^T)$ for all $x, y \in V$ $(x \neq y)$, and

(b) for each $e = xy \in E(T)$, the elementary cocycle C of G corresponding to e and T is a minimum-weight edge cut separating x from y in G, i.e. $\lambda_G(x, y; \mathbf{w}) = \sum\limits_{f \in C} w_f$.

We shall call any tree T with the above properties a *Gomory–Hu tree* for $(G, \mathbf{w})$; it is in general nonunique. Note that, for any given tree T, there is at most one choice of $\mathbf{w}^T$ that satisfies (a), namely for each edge $e = xy \in E(T)$ we must set $w_e^T = \lambda_G(x, y; \mathbf{w})$. It can also be shown that if T satisfies (b), then this definition of $\mathbf{w}^T$ necessarily satisfies (a); but we shall not need this fact.

If T is a Gomory–Hu tree for $(G, \mathbf{w})$, we define $\widehat{\Lambda}(G, \mathbf{w}; T) = \max_{e \in E(T)} w_e^T$. We claim that this value is independent of the choice of T, and in fact we have:

Theorem 2.8 *Let G be a connected graph equipped with nonnegative real edge weights $\mathbf{w} = \{w_e\}_{e \in E(G)}$, and let T be a Gomory–Hu tree for $(G, \mathbf{w})$. Then*

$$\Lambda(G, \mathbf{w}) \;=\; \widehat{\Lambda}(G, \mathbf{w}; T) \;=\; \widetilde{\Lambda}(G, \mathbf{w}) \;\le\; \Delta_2(G, \mathbf{w}) \;\le\; \Delta(G, \mathbf{w}) \;. \tag{2.1}$$

In particular, the value of $\widehat{\Lambda}(G, \mathbf{w}; T)$ is independent of the choice of T.

Proof. The equality $\Lambda(G, \mathbf{w}) = \Lambda(T, \mathbf{w}^T)$ follows from Theorem 2.7(a), and it is trivial to see that $\Lambda(T, \mathbf{w}^T) = \max_{e \in E(T)} w_e^T$. This proves that $\Lambda(G, \mathbf{w}) = \widehat{\Lambda}(G, \mathbf{w}; T)$ and in particular that the latter quantity is independent of the choice of T.

The inequality $\Lambda(G, \mathbf{w}) \le \widetilde{\Lambda}(G, \mathbf{w})$ follows from Lemma 2.6.

The inequality $\widetilde{\Lambda}(G, \mathbf{w}) \le \widehat{\Lambda}(G, \mathbf{w}; T)$ follows from Lemma 2.5 and Theorem 2.7(b).

There are easy elementary proofs of both $\Lambda(G, \mathbf{w}) \le \Delta_2(G, \mathbf{w})$ and $\widetilde{\Lambda}(G, \mathbf{w}) \le \Delta_2(G, \mathbf{w})$, as noted in the Introduction. □

Corollary 2.9 *Let G be a (not necessarily connected) graph equipped with nonnegative real edge weights $\mathbf{w} = \{w_e\}_{e \in E(G)}$. Then*

$$\Lambda(G, \mathbf{w}) \;=\; \widetilde{\Lambda}(G, \mathbf{w}) \;\le\; \Delta_2(G, \mathbf{w}) \;\le\; \Delta(G, \mathbf{w}) \;. \tag{2.2}$$

Proof. If G is disconnected, it suffices to apply Theorem 2.8 to each component of G. □

Finally, we need to prove our claims that $\Lambda(G, \mathbf{w}) \ge D(G, \mathbf{w})$ and $\Lambda(G, \mathbf{w}) \ge \Delta_{n-1}(G, \mathbf{w})$. We shall actually prove a slightly stronger result. Define the k^{th} *weighted degeneracy number*

$$D_k(G, \mathbf{w}) \;=\; \max_{H \subseteq G} \delta_k(H, \mathbf{w}) \;, \tag{2.3}$$

where the max runs over all subgraphs H of G, and $\delta_k(H, \mathbf{w})$ denotes the kth smallest weighted degree of H:

$$\delta_k(H, \mathbf{w}) \;=\; \max_{x_1,\ldots,x_{k-1} \in V(H)} \; \min_{x \in V(H) \setminus \{x_1,\ldots,x_{k-1}\}} d_H(x, \mathbf{w}) \;. \tag{2.4}$$

Trivially we have $D(G, \mathbf{w}) \equiv D_1(G, \mathbf{w}) \le D_2(G, \mathbf{w}) \le \ldots$ and $\delta_k(G, \mathbf{w}) \le D_k(G, \mathbf{w})$. In particular,

$$D_2(G, \mathbf{w}) \;\ge\; \max[D(G, \mathbf{w}),\, \delta_2(G, \mathbf{w})] \;. \tag{2.5}$$

Proposition 2.10 *Let G be a graph with n vertices $(n \ge 2)$ equipped with nonnegative real edge weights $\mathbf{w} = \{w_e\}_{e \in E(G)}$. Then*

$$\Lambda(G, \mathbf{w}) \;\ge\; D_2(G, \mathbf{w}) \;\ge\; \max[D(G, \mathbf{w}),\, \Delta_{n-1}(G, \mathbf{w})] \;. \tag{2.6}$$

PROOF. Suppose first that G is connected, and let T be a Gomory–Hu tree for $(G, \mathbf{w})$. For any vertex x of degree 1 in T, let $e = xy$ be the unique incident edge in T; then the elementary cocycle of G corresponding to e and T is $E_G(\{x\}, \{x\}^c)$. Using Theorem 2.7(b), we have $\Lambda(G, \mathbf{w}) \ge \lambda_G(x, y; \mathbf{w}) = d_G(x, \mathbf{w})$. Since there are at least two such vertices x, we have $\Lambda(G, \mathbf{w}) \ge \delta_2(G, \mathbf{w})$.

If G is disconnected, we can apply the result just proven to each component of G; we conclude again that $\Lambda(G, \mathbf{w}) \ge \delta_2(G, \mathbf{w})$.

Now apply this result to each subgraph H of G: we conclude that $\Lambda(H, \mathbf{w}|_H) \ge \delta_2(H, \mathbf{w})$. But $\Lambda(G, \mathbf{w}) \ge \Lambda(H, \mathbf{w}|_H)$ for every subgraph H of G, so $\Lambda(G, \mathbf{w}) \ge D_2(G, \mathbf{w})$. $\square$

Let us now prove a few further general properties of maxmaxflow. Let G be a graph and $x \in V(G)$. We say that x is a *cut vertex* of G if $G \setminus x$ has more components than G. We say that G is *non-separable* if G is connected and has no cut vertices. A *block* of G is a maximal non-separable subgraph of G. We first observe that maxmaxflow has a naturalness property that maximum degree and kth-largest degree lack, namely, it "trivializes over blocks":

Proposition 2.11 *Let $G_1, \ldots, G_b$ be the blocks of G. Then $\Lambda(G, \mathbf{w}) = \max\limits_{1 \le i \le b} \Lambda(G_i, \mathbf{w})$.*

PROOF. If x and y lie in the same block G_i, then $\lambda_G(x, y; \mathbf{w}) = \lambda_{G_i}(x, y; \mathbf{w})$. If x and y lie in the same component of G but in different blocks, then there exist cut vertices $v_1, \ldots, v_k$ of G and blocks $G_{i_0}, G_{i_1}, \ldots, G_{i_k}$ of G such that $x \in V(G_{i_0})$, $y \in V(G_{i_k})$, $V(G_{i_{j-1}}) \cap V(G_{i_j}) = \{v_j\}$, and every path from x to y passes through $v_1, \ldots, v_k$ in that order; and in this case we have

$$\lambda_G(x, y; \mathbf{w}) \;=\; \min\Bigl[\lambda_{G_0}(x, v_1; \mathbf{w}),\, \lambda_{G_1}(v_1, v_2; \mathbf{w}),\, \ldots,\, \lambda_{G_{k-1}}(v_{k-1}, v_k; \mathbf{w}),\, \lambda_{G_k}(v_k, y; \mathbf{w})\Bigr] \;. \tag{2.7}$$

Finally, if x and y lie in different components of G, then $\lambda_G(x, y; \mathbf{w}) = 0$. $\square$

It follows immediately from the definition of maxmaxflow that for any pair of distinct vertices x_1, x_2 of G, there exists a partition $V(G) = X_1 \cup X_2$ such that $x_1 \in X_1$, $x_2 \in X_2$ and $\sum_{e \in E(X_1, X_2)} w_e = \lambda_G(x_1, x_2; \mathbf{w}) \leq \Lambda(G, \mathbf{w})$. We will need the following extension of this observation in Section 7.

Proposition 2.12 *Let* $X \subseteq V(G)$ *with* $|X| \geq 2$. *Then there exist* $x_1, x_2 \in X$ *and disjoint* $X_1, X_2 \subseteq V(G)$ *such that* $X \cap X_i = \{x_i\}$ *and* $\sum_{e \in E(X_i, X_i^c)} w_e \leq \Lambda(G, \mathbf{w})$ *for all* $1 \leq i \leq 2$.

Proof. We can assume without loss of generality that G is connected. Let T be a Gomory–Hu tree for $(G, \mathbf{w})$, and let T' be the union of all the paths in T connecting pairs of vertices of X. Let x_1, x_2 be distinct end-vertices of T' and e_1, e_2 be the edges of T' incident with x_1, x_2 respectively. For $1 \leq i \leq 2$, let X_i to be the vertex set of the component of $T \setminus e_i$ which contains x_i. Then $X_1 \cap X_2 = \varnothing$. Furthermore, $E(X_i, X_i^c)$ is the elementary cocycle of G corresponding to the edge e_i of T; so by Theorem 2.7(b) we have $\sum_{e \in E(X_i, X_i^c)} w_e \leq \Lambda(G, \mathbf{w})$ for all $1 \leq i \leq 2$. $\square$

We conclude this section with a few examples of maxmaxflow calculations:

Example 2.1. Let G be any forest, and let $\mathbf{w}$ be any set of nonnegative edge weights. Then $\Lambda(G, \mathbf{w}) = \max_{e \in E(G)} w_e$. Indeed, this elementary fact was already used in the proof of Theorem 2.8. $\square$

Example 2.2. Let G be the cycle C_n, and let $\mathbf{w}$ be any set of nonnegative edge weights. Then

$$\Lambda(G, \mathbf{w}) \;=\; \max_{e \in E(G)} w_e \,+\, \min_{e \in E(G)} w_e \,. \tag{2.8}$$

Proof. For any pair of distinct vertices $x, y \in V(G)$, there are precisely two paths from x to y, and together they use all the edges of G. Hence on one path the max flow is exactly $\min w_e$, and on the other it is at most $\max w_e$. So $\Lambda(G, \mathbf{w}) \leq \min w_e + \max w_e$. On the other hand, if we take x, y to be the endpoints of the edge with maximum weight, we obtain equality. $\square$

3 Some Preliminaries

Before turning to the counting of subgraphs of various classes, let us make some brief prefatory observations.

3.1 Pointwise bounds vs. generating-function bounds

Let $(G, \mathbf{w})$ be a weighted graph, and suppose that a_m is the total weight of all m-edge subgraphs of G of some specified class. In the following sections we shall prove upper bounds on a_m of two different types:

- "Pointwise bound": $a_m \le C_m$ for some specified constants C_m.
- "Generating-function bound": $\sum_{m=0}^{\infty} a_m z^m \le C(z)$ for some specified function $C(z)$ [which is allowed to take the value $+\infty$], for all $z \ge 0$.

From each type of bound we can deduce one of the other type: $a_m \le C_m$ trivially implies $\sum a_m z^m \le \sum C_m z^m$; and $\sum a_m z^m \le C(z)$ implies $a_m \le \inf_{z \ge 0} z^{-m} C(z)$. But we lose something in each such passage: for example,

$$\sum_{m=0}^{\infty} a_m \le 1 \implies a_m \le 1 \implies \sum_{m=0}^{\infty} a_m z^m \le \frac{1}{1-z} \implies a_m \le \frac{(m+1)^{m+1}}{m^m} \implies \cdots \tag{3.1}$$

So it is important to choose the best type of bound in each problem. Roughly speaking, the pointwise bound is better in those cases where the bound can be saturated simultaneously for all (or arbitrarily many) values of m. By contrast, in cases where the pointwise bound can be saturated only for one value of m at a time, it may be possible to prove a stronger bound of generating-function type. These remarks will be illustrated repeatedly throughout the following sections.

3.2 Convex hulls in graphs

Let H be any graph, and let X be a nonempty subset of $V(H)$. We define the *convex hull of X in H*, denoted $\operatorname{conv}(X, H)$, to be the union of all paths in H connecting any pair of vertices $x_1, x_2 \in X$ (including paths of length 0 from a vertex $x \in X$ to itself). Thus, $\operatorname{conv}(X, H)$ is a subgraph of H whose vertex set contains X. The following properties are elementary consequences of this definition:

(Conv1) If H is connected, then $\operatorname{conv}(X, H)$ is connected.

(Conv2) Any vertex of degree 0 or 1 in $\operatorname{conv}(X, H)$ must belong to X. Moreover, a vertex $x \in X$ is of degree 0 (i.e. isolated) in $\operatorname{conv}(X, H)$ if and only if it is the only element of X in its component of H.

(Conv3) If H is a tree, then $\operatorname{conv}(X, H)$ is the smallest subtree of H containing all the vertices of X.

(Conv4) If H is 2-connected (or isomorphic to K_2) and $|X| \ge 2$, then $\operatorname{conv}(X, H) = H$. [This is because, for any pair of distinct vertices x_1, x_2 of H, every edge of H lies on some path from x_1 to x_2.]

Property (Conv4) can be generalized as follows. Let H be a graph and $x \in V(H)$. We say that x is an *internal vertex* of H if x is not a cut vertex of H. An *end block* of H is a block that contains exactly one cut vertex of H.

Lemma 3.1 *Let H be connected, and let $X \subseteq V(H)$ with $|X| \geq 2$. Then:*

(a) $\operatorname{conv}(X, H)$ *is a connected union of blocks of H.*

(b) Each end block of $\operatorname{conv}(X, H)$ *has an internal vertex belonging to X.*

PROOF. This follows easily from the definition of $\operatorname{conv}(X, H)$ and property (Conv4). $\square$

We have already used convex hulls in the proof of Proposition 2.12, and they will play an important role in our treatment of trees and block trees (Sections 5 and 7).

4 Counting Walks and Paths

Let G be a graph equipped with nonnegative real edge weights $\mathbf{w} = \{w_e\}_{e \in E(G)}$. In this section (as well as in the following ones) we shall write Δ, Δ_2, Λ, ... as a shorthand for $\Delta(G, \mathbf{w})$, $\Delta_2(G, \mathbf{w})$, $\Lambda(G, \mathbf{w})$, ...; the underlying graph G and its edge weights $\mathbf{w}$ will always be understood. Similarly, we shall write $\lambda(x, y)$ as a shorthand for $\lambda_G(x, y; \mathbf{w})$.

For $x, y \in V(G)$ and $m \geq 0$, let $\mathcal{W}_m(x, y)$ be the set of m-step walks from x to y, i.e. sequences $\omega = x_0 e_1 x_1 e_2 x_2 \cdots x_{m-1} e_m x_m$ with $x_0 = x$ and $x_m = y$ such that each e_i is an edge $x_{i-1} x_i$. We then define the following subsets of $\mathcal{W}_m(x, y)$:

- $\mathcal{W}_m^{\mathrm{FPW}}(x, y)$ consists of *first-passage walks*, i.e. walks such that $x_i \neq y$ for $i < m$;
- $\mathcal{W}_m^{\mathrm{SAW}}(x, y)$ consists of *self-avoiding walks* (otherwise known as *paths*), i.e. walks such that $x_i \neq x_j$ for $i \neq j$. [We remark that a walk ω is self-avoiding if and only if each initial segment of ω is a first-passage walk.]

Clearly $\mathcal{W}_m^{\mathrm{SAW}}(x, y) \subseteq \mathcal{W}_m^{\mathrm{FPW}}(x, y) \subseteq \mathcal{W}_m(x, y)$.

More generally, let $x \in V(G)$ and $Y \subseteq V(G)$. We let $\mathcal{W}_m(x, Y) = \bigcup_{y \in Y} \mathcal{W}_m(x, y)$ be the set of m-step walks from x to Y. We then define the following subsets of $\mathcal{W}_m(x, Y)$:

- $\mathcal{W}_m^{\mathrm{FPW}}(x, Y)$ consists of *first-passage walks to the set Y*, i.e. walks such that $x_i \notin Y$ for $i < m$;
- $\mathcal{W}_m^{\mathrm{FPSAW}}(x, Y)$ consists of *first-passage self-avoiding walks to the set Y*, i.e. walks such that $x_i \notin Y$ for $i < m$ and $x_i \neq x_j$ for $i \neq j$.

Then $\mathcal{W}_m^{\rm FPSAW}(x,Y) \subseteq \mathcal{W}_m^{\rm FPW}(x,Y) \subseteq \mathcal{W}_m(x,Y)$, and in fact $\mathcal{W}_m^{\rm FPSAW}(x,Y) = \mathcal{W}_m^{\rm FPW}(x,Y) \cap \bigcup\limits_{y \in Y} \mathcal{W}_m^{\rm SAW}(x,y)$.

We define the *weight* of a walk $\omega = x_0 e_1 x_1 e_2 x_2 \cdots x_{m-1} e_m x_m$ to be the product of its edge weights:

$$w(x_0 e_1 x_1 e_2 x_2 \cdots x_{m-1} e_m x_m) \;=\; \prod_{i=1}^{m} w_{e_i} \;. \tag{4.1}$$

Finally, we define the weighted counts

$$w_m(x,y) \;=\; \sum_{\omega \in \mathcal{W}_m(x,y)} w(\omega) \tag{4.2}$$

$$w_m(x,Y) \;=\; \sum_{\omega \in \mathcal{W}_m(x,Y)} w(\omega) \tag{4.3}$$

and likewise for $w^{\rm FPW}$, $w^{\rm SAW}$ and $w^{\rm FPSAW}$.

Walks with a fixed initial vertex and an arbitrary final vertex can trivially be bounded in terms of maximum weighted degree:

Proposition 4.1 *For all $x \in V(G)$ and all $m \geq 0$, we have*

$$\sum_{y \in V(G)} w_m(x,y) \;\leq\; \Delta^m \;. \tag{4.4}$$

In fact, when $(G, \mathbf{w})$ is Δ-regular [i.e. $d_G(x, \mathbf{w}) = \Delta$ for all $x \in V(G)$], the bound (4.4) is sharp simultaneously for all $x \in V(G)$ and all $m \geq 0$.

If we restrict to first-passage walks to a fixed target set Y, we can obtain a "generating-function" bound that is sharper than (4.4):

Proposition 4.2 *For all $x \in V(G)$ and all $Y \subseteq V(G)$, we have*

$$\sum_{m=0}^{\infty} \Delta^{-m} w_m^{\rm FPW}(x,Y) \;\leq\; 1 \;. \tag{4.5}$$

PROOF. Define a sub-Markov chain with transition probabilities $p(x \to x') = \Delta^{-1} \sum_{e = xx'} w_e$ (here the sum runs over all edges $e = xx'$ in G). Then $\sum_{m=0}^{\infty} \Delta^{-m} w_m^{\rm FPW}(x,Y)$ is the probability that Y is eventually hit, starting at x. $\square$

Remarks. 1. If G is connected (with weights $w_e > 0$) and $(G, \mathbf{w})$ is Δ-regular, then (4.5) is *equality* for all $x \in V(G)$ and all $Y \subseteq V(G)$. This is because p is an ergodic Markov chain, so that Y is hit with probability 1. Indeed, it suffices to have $d_G(x, \mathbf{w}) = \Delta$ for all $x \in V(G) \setminus Y$; the degree at vertices of Y is irrelevant.

2. If $x \in Y$ then $w_0^{\rm FPW}(x,Y) = 1$ and $w_m^{\rm FPW}(x,Y) = 0$ for $m \geq 1$ so equality holds in (4.5). On the other hand, if $x \notin Y$, then we can improve the upper bound in (4.5) from 1 to $d_G(x,\mathbf{w})/\Delta$, since this is the probability for the Markov chain to survive the first step.

3. As just mentioned, (4.5) is the best-possible upper bound for first-passage walks to a set Y. However, one might ask whether a sharper bound is possible by restricting attention to first-passage *self-avoiding* walks. The answer is no, at least if one considers a general set Y, as the following examples show:

Example 4.1. Let G be the star $K_{1,r}$, let x be the central vertex, and let Y be the remaining vertices. Set all edge weights $w_e = \Delta/r$. Then $w_1^{\rm FPSAW}(x,Y) = \Delta$ and $w_m^{\rm FPSAW}(x,Y) = 0$ for $m \neq 1$, so (4.5) is sharp. $\square$

Example 4.2. More generally, let $\mathbf{T}_r$ be the infinite r-regular tree, let x be any vertex of $\mathbf{T}_r$, and fix $n \geq 1$. Let G be the subtree of $\mathbf{T}_r$ induced by the vertices lying at a distance at most n from x, and let Y be the set of vertices lying at a distance exactly n from x. Set all edge weights $w_e = \Delta/r$. Then

$$w_m^{\rm FPSAW}(x,Y) \;=\; \begin{cases} r(r-1)^{m-1}(\Delta/r)^m & \text{if } m=n \\ 0 & \text{otherwise} \end{cases} \tag{4.6}$$

so that

$$\sum_{m=0}^{\infty} \Delta^{-m} w_m^{\rm FPSAW}(x,Y) \;=\; \left(\frac{r-1}{r}\right)^{n-1} \;, \tag{4.7}$$

which is sharp in the limit $r \to \infty$ at any fixed n. (It is also sharp for any r when $n = 1$, which is the case $G = K_{1,r}$.) $\square$

For comparison with Proposition 4.3 below, it is perhaps illuminating to rephrase the proof of Proposition 4.2 as an induction:

Second proof of Proposition 4.2. We will prove inductively that

$$\sum_{m=0}^{M} \Delta^{-m} w_m^{\rm FPW}(x,Y) \leq 1$$

for all $M \geq 0$ and all $x \in V(G)$. This clearly holds for $M = 0$. Moreover, it holds for all M when $x \in Y$, since $w_m^{\rm FPW}(x,Y) = \delta_{m0}$. So suppose that $x \notin Y$. Then $w_0^{\rm FPW}(x,Y) = 0$ and for $m \geq 1$ we have

$$w_m^{\rm FPW}(x,Y) \;=\; \sum_{x' \in V(G)} \left(\sum_{e=xx'} w_e \right) w_{m-1}^{\rm FPW}(x',Y) \;. \tag{4.8}$$

Thus, for $M \geq 1$, we have

$$\begin{aligned} \sum_{m=0}^{M} \Delta^{-m} w_m^{\mathrm{FPW}}(x,Y) &= \sum_{m=1}^{M} \Delta^{-m} \sum_{x' \in V(G)} \left(\sum_{e=xx'} w_e \right) w_{m-1}^{\mathrm{FPW}}(x',Y) \\ &= \Delta^{-1} \sum_{x' \in V(G)} \left(\sum_{e=xx'} w_e \right) \sum_{j=0}^{M-1} \Delta^{-j} w_j^{\mathrm{FPW}}(x',Y) \\ &\leq 1 \quad \text{by the inductive hypothesis.} \end{aligned} \tag{4.9}$$

□

There is no hope of obtaining a $|Y|$-independent bound on $w_m^{\mathrm{FPW}}(x,Y)$ — or even on $w_m^{\mathrm{FPSAW}}(x,Y)$ — in terms of second-largest degree (much less in terms of maxmaxflow). Indeed, let G be the star $K_{1,r}$, so that $\Delta = r$ and $\Delta_2 = \Lambda = 1$; let x be the central vertex and Y the remaining vertices; then $w_1^{\mathrm{FPSAW}}(x,Y) = r$, which is unbounded. Nor can we bound $w_m^{\mathrm{FPW}}(x,y)$: considering again the star and taking y to be any vertex other than x, we have $w_3^{\mathrm{FPW}}(x,y) = r-1$, which is again unbounded. Nevertheless, an analogue of Proposition 4.2 does hold for maxmaxflow if we restrict ourselves to self-avoiding walks and to the case $Y = \{y\}$:

Proposition 4.3 *Define*

$$F(x,y) \;=\; \begin{cases} \lambda(x,y)/\Lambda & \text{if } x \neq y \\ 1 & \text{if } x = y \end{cases} \tag{4.10}$$

Then, for all $x, y \in V(G)$, we have

$$\sum_{m=0}^{\infty} \Lambda^{-m} w_m^{\mathrm{SAW}}(x,y) \;\leq\; F(x,y) \;\leq\; 1 \;. \tag{4.11}$$

PROOF. We will prove inductively that $\sum_{m=0}^{M} \Lambda^{-m} w_m^{\mathrm{SAW}}(x,y) \leq F(x,y)$ for all $M \geq 0$ and all $x, y \in V(G)$. This clearly holds for $M = 0$, since $w_0^{\mathrm{SAW}}(x,y) = \delta_{xy}$. Moreover, it holds for all M when $x = y$, since $w_m^{\mathrm{SAW}}(x,x) = \delta_{m0}$. So let $M \geq 1$ and $x \neq y$. Let $C = E(X,Y)$ be a cocycle in G with $x \in X$, $y \in Y$ and $\sum_{e \in C} w_e = \lambda(x,y)$. Let $e = uv$ be an edge in C with $u \in X$ and $v \in Y$, and let $w_m^{\mathrm{SAW}}(x,y,e)$ be the weighted sum over m-step paths from x to y that use e as their first edge from X to Y. Then

$$w_m^{\mathrm{SAW}}(x,y,e) \;\leq\; w_e \sum_{i=0}^{m-1} w_i^{\mathrm{SAW}}(x,u) w_{m-1-i}^{\mathrm{SAW}}(v,y) \;. \tag{4.12}$$

So

$$\begin{aligned}\sum_{m=0}^{M} \Lambda^{-m} w_m^{\mathrm{SAW}}(x,y,e) &\le \Lambda^{-1} w_e \sum_{m=0}^{M} \sum_{i=0}^{m-1} \Lambda^{-i} w_i^{\mathrm{SAW}}(x,u) \Lambda^{-(m-1-i)} w_{m-1-i}^{\mathrm{SAW}}(v,y) \\ &= \Lambda^{-1} w_e \sum_{i=0}^{M-1} \Lambda^{-i} w_i^{\mathrm{SAW}}(x,u) \sum_{j=0}^{M-1-i} \Lambda^{-j} w_j^{\mathrm{SAW}}(v,y) \\ &\le \Lambda^{-1} w_e \quad \text{by the inductive hypothesis} \qquad (4.13)\end{aligned}$$

(note that we may have here used the $x = y$ case of the inductive hypothesis, in case $x = u$ or $y = v$ or both). Therefore

$$\begin{aligned}\sum_{m=0}^{M} \Lambda^{-m} w_m^{\mathrm{SAW}}(x,y) &= \sum_{e\in C} \sum_{m=0}^{M} \Lambda^{-m} w_m^{\mathrm{SAW}}(x,y,e) \\ &\le \sum_{e\in C} \Lambda^{-1} w_e \\ &\le \lambda(x,y)/\Lambda\,. \qquad (4.14)\end{aligned}$$

□

Corollary 4.4 *$w_m^{\mathrm{SAW}}(x,y) \le \Lambda^m F(x,y)$ for all $m \ge 0$ and all $x, y \in V(G)$.*

Corollary 4.5 *For $0 \le \zeta \le \Lambda^{-1}$, we have $\sum\limits_{m=0}^{\infty} \zeta^m w_m^{\mathrm{SAW}}(x,y) \le (\zeta\Lambda)^{\mathrm{dist}(x,y)} F(x,y)$ where $\mathrm{dist}(x,y)$ is the length of the shortest path in G from x to y. (Of course, if there is no such path, then $w_m^{\mathrm{SAW}}(x,y) = 0$ for all m.)*

Example 4.3. Let G be any tree, let $w_e = w$ for all edges e, and let x, y be any two vertices of G. Then $\lambda(x,y) = \Lambda = w$ and

$$w_m^{\mathrm{SAW}}(x,y) \;=\; \begin{cases} w^m & \text{if } m = \mathrm{dist}(x,y) \\ 0 & \text{otherwise} \end{cases} \qquad (4.15)$$

so that (4.11) is equality for all x, y. Note that the generating-function bound (4.11) is here much sharper than the pointwise bound of Corollary 4.4: this reflects the fact that the latter can be saturated here for only one value of m (for any given pair x, y). This is the prototypical situation in which we shall seek generating-function bounds. Indeed, the inductive “cutting” argument used in the proof of Proposition 4.3 seems to work *only* for the generating-function bound; we do not know of any way of proving Corollary 4.4 without proving the stronger bound (4.11). ☐

Example 4.4. Let G be the complete graph K_n, and let all edge weights w_e equal $\Delta/(n-1)$, so that both the maximum weighted degree and the maxmaxflow equal Δ. If x, y are any two distinct vertices, we have

$$w_m^{\rm SAW}(x,y) \;=\; \frac{(n-2)(n-3)\cdots(n-m)}{(n-1)^m}\,\Delta^m \tag{4.16}$$

if $1 \le m \le n-1$, and 0 otherwise. Approximating Riemann sums by integrals, we find

$$\sum_{m=0}^{\infty} \Delta^{-m} w_m^{\rm SAW}(x,y) \;\approx\; \sqrt{\frac{\pi}{2n}} \tag{4.17}$$

as $n \to \infty$, so Proposition 4.3 is far from sharp in this limit. Nevertheless, we see that the exponential growth rate $w_m^{\rm SAW}(x,y) \sim \Delta^m$ in Proposition 4.3 and Corollary 4.4 cannot be improved. Indeed, if for each fixed m we choose n so as to maximize $w_m^{\rm SAW}(x,y)/\Delta^m$, we find that the maximum is achieved at $n \approx m^2/2$ and that the maximum value is $\approx e/(2m^2)$. ☐

Example 4.5. Let G be the generalized theta graph $\Theta_{1,2,\ldots,r}$ ($r \ge 2$), consisting of a pair of endvertices a, b joined by r internally disjoint paths of lengths $1, 2, \ldots, r$. On each path let one edge have weight w and the other edges have weight 1. Then the maxmaxflow is

$$\Lambda \;=\; \begin{cases} 1+w & \text{if } 0 \le w \le 1/(r-1) \\ rw & \text{if } 1/(r-1) \le w \le 1 \\ r-1+w & \text{if } w \ge 1 \end{cases} \tag{4.18}$$

while

$$w_m^{\rm SAW}(a,b) \;=\; \begin{cases} w & \text{if } 1 \le m \le r \\ 0 & \text{otherwise} \end{cases} \tag{4.19}$$

In particular, at $w = 1/(r-1)$ we have

$$\sum_{m=0}^{\infty} \Lambda^{-m} w_m^{\rm SAW}(a,b) \;=\; 1 \,-\, \left(1 - \frac{1}{r}\right)^{r} \;, \tag{4.20}$$

which decreases to $1 - 1/e \approx 0.632121$ as $r \to \infty$. So, although the bound of Proposition 4.3 is not sharp in this case, it does at least come within a constant factor of being sharp in a situation where the maximum contribution from a single value of m goes to zero (the opposite extreme from Examples 4.1 and 4.2). ☐

5 Counting Trees and Forests

Let us now extend Propositions 4.2 and 4.3 from paths to trees and forests. In Section 5.1 we consider classes $\mathcal{T}_m(X)$ of trees and $\mathcal{F}_m(X,Y)$ of forests. In Section 5.2 we consider a larger class $\mathcal{H}_m(X)$ of forests.

5.1 The classes $\mathcal{T}_m(X)$ and $\mathcal{F}_m(X,Y)$

For F a forest in G, let $L(F)$ denote the set of vertices of degree 0 or 1 in F (also called *leaves* or *end-vertices* of F).

For any nonempty $X \subseteq V(G)$, let $\mathcal{T}_m(X)$ be the set of all m-edge trees T in G such that $L(T) \subseteq X \subseteq V(T)$. Heuristically, $\mathcal{T}_m(X)$ consists of trees whose leaves are "tied down" on the set X. Note the following special cases:

- If $X = \{x\}$, then $\mathcal{T}_0(\{x\})$ has as its single element the edgeless graph with vertex set $\{x\}$, and $\mathcal{T}_m(\{x\}) = \varnothing$ for $m \geq 1$.
- $\mathcal{T}_m(\{x,y\}) \simeq \mathcal{W}_m^{\mathrm{SAW}}(x,y)$ under the natural identification of paths with their induced subgraphs; this holds both for $x \neq y$ and for $x = y$.
- $\mathcal{T}_0(X) = \varnothing$ for $|X| \geq 2$.

For any $X, Y \subseteq V(G)$ with $Y \neq \varnothing$, let $\mathcal{F}_m(X,Y)$ be the set of all m-edge forests F in G such that

(F1) $L(F) \subseteq X \cup Y \subseteq V(F)$, and

(F2) each component of F contains exactly one vertex of Y.

[Note that $\mathcal{F}_m(X,Y) = \mathcal{F}_m(X \setminus Y, Y)$, so we can assume without loss of generality, if desired, that $X \cap Y = \varnothing$.] Heuristically, $\mathcal{F}_m(X,Y)$ consists of forests whose leaves are "tied down" on the set X and whose components are "tied down" on single elements of the set Y. We have the following special cases:

- If $X = \varnothing$ (or more generally if $X \subseteq Y$), then $\mathcal{F}_0(X,Y)$ has as its single element the edgeless graph with vertex set Y, and $\mathcal{F}_m(X,Y) = \varnothing$ for $m \geq 1$.
- If $X = \{x\}$, then each $F \in \mathcal{F}_m(\{x\},Y)$ is the disjoint union of a path $P \in \mathcal{W}_m^{\mathrm{FPSAW}}(x,Y)$ [under the natural identification of paths with their induced subgraphs] and the collection $Y \setminus V(P)$ of isolated vertices; this holds both for $x \notin Y$ and for $x \in Y$. We express this isomorphism loosely by writing $\mathcal{F}_m(\{x\},Y) \simeq \mathcal{W}_m^{\mathrm{FPSAW}}(x,Y)$.
- For $Y = \{y\}$, we have $\mathcal{F}_m(X,\{y\}) = \mathcal{T}_m(X \cup \{y\})$.

For H a subgraph of G, set $w(H) = \prod_{e \in E(H)} w_e$. [Note that if $E(H) = \varnothing$, then $w(H) = 1$.] Define the weighted counts

$$t_m(X) \;=\; \sum_{T \in \mathcal{T}_m(X)} w(T) \tag{5.1}$$

$$f_m(X, Y) \;=\; \sum_{F \in \mathcal{F}_m(X,Y)} w(F) \tag{5.2}$$

We will obtain two "generating-function" bounds on $f_m(X, Y)$: a sharp bound in terms of the maximum weighted degree Δ, and a slightly weaker bound in terms of the maxmaxflow Λ.

Proposition 5.1 *For all $X, Y \subseteq V(G)$ with $Y \neq \varnothing$, we have*

$$\sum_{m=0}^{\infty} \Delta^{-m} f_m(X, Y) \;\le\; 1 \,. \tag{5.3}$$

Proposition 5.2 *For all $X, Y \subseteq V(G)$ with $Y \neq \varnothing$, we have*

$$\sum_{m=0}^{\infty} (m + |Y|)^{-(|X|-1)} \Lambda^{-m} f_m(X, Y) \;\le\; |Y| \,. \tag{5.4}$$

Using the identity $\mathcal{F}_m(X, \{y\}) = \mathcal{T}_m(X \cup \{y\})$, we immediately obtain the following corollaries for trees:

Corollary 5.3 *For all nonempty $X \subseteq V(G)$, we have*

$$\sum_{m=0}^{\infty} \Delta^{-m} t_m(X) \;\le\; 1 \,. \tag{5.5}$$

Corollary 5.4 *For all nonempty $X \subseteq V(G)$, we have*

$$\sum_{m=0}^{\infty} (m + 1)^{-(|X|-2)} \Lambda^{-m} t_m(X) \;\le\; 1 \,. \tag{5.6}$$

One can envisage two different approaches to proving these bounds for forests and trees. One is to *imitate* the "cutting" argument employed for walks in the proofs of Propositions 4.2 and 4.3. The other is to *exploit the result* obtained for walks in Propositions 4.2 and 4.3, by reducing forests inductively to paths. The latter technique seems to be more appropriate.

In the proof of Proposition 5.1 we shall make use of the "point-to-set" bound of Proposition 4.2. We shall also use the following elementary lemma which splits a forest with k end-vertices into a forest with $k - 1$ end-vertices and a path:

Lemma 5.5 *Let G be a graph, let $X, Y \subseteq V(G)$ with $Y \neq \varnothing$, let $x \in X \setminus Y$, and let $F \in \mathcal{F}_m(X,Y)$. Let F_1 be the convex hull of $(X \setminus x) \cup Y$ in F, and let P be the unique path in F from x to $V(F_1)$. Then F is the edge-disjoint union of F_1 and P; and for some i $(0 \le i \le m)$ we have $F_1 \in \mathcal{F}_i(X \setminus x, Y)$ and $P \in \mathcal{W}_{m-i}^{\rm FPSAW}(x, V(F_1))$. Moreover, the map $F \mapsto (F_1, P)$ is an injection.*

The proof is a straightforward exercise: let us simply observe that because $x \notin Y$, the component of F containing x must contain a vertex in Y, which guarantees that the path P exists; and P is unique because F has no cycles.

Proof of Proposition 5.1. As noted above, we can assume without loss of generality that $X \cap Y = \varnothing$. We use induction on $k = |X|$. For $k = 0$ the result is trivial. For $k = 1$ the result follows immediately from Proposition 4.2, since $\mathcal{F}_m(\{x\}, Y) \simeq \mathcal{W}_m^{\rm FPSAW}(x,Y) \subseteq \mathcal{W}_m^{\rm FPW}(x,Y)$. So suppose $k \ge 2$. Let x be any vertex in X; by assumption $x \notin Y$. By Lemma 5.5, given any $F \in \mathcal{F}_m(X,Y)$, we can decompose F into a forest $F_1 \in \mathcal{F}_i(X \setminus x, Y)$ and a path $P \in \mathcal{W}_{m-i}^{\rm FPSAW}(x, V(F_1))$; and each such pair (F_1, P) arises from a unique F (namely, $F_1 \cup P$). Since $w(F) = w(F_1)w(P)$, we have

$$
\begin{aligned}
&\sum_{m=0}^{\infty} \Delta^{-m} \sum_{F \in \mathcal{F}_m(X,Y)} w(F) \\
&\quad \le \sum_{m=0}^{\infty} \sum_{i=0}^{m} \Delta^{-i} \sum_{F_1 \in \mathcal{F}_i(X \setminus x, Y)} w(F_1)\, \Delta^{-(m-i)} \sum_{P \in \mathcal{W}_{m-i}^{\rm FPSAW}(x, V(F_1))} w(P) \\
&\quad = \sum_{i=0}^{\infty} \Delta^{-i} \sum_{F_1 \in \mathcal{F}_i(X \setminus x, Y)} w(F_1) \sum_{j=0}^{\infty} \Delta^{-j} \sum_{P \in \mathcal{W}_{j}^{\rm FPSAW}(x, V(F_1))} w(P)\,. \qquad (5.7)
\end{aligned}
$$

Now, for each fixed F_1 we have $\sum_{j=0}^{\infty} \Delta^{-j} \sum_{P \in \mathcal{W}_j^{\rm FPSAW}(x, V(F_1))} w(P) \le 1$ by Proposition 4.2, so that

$$
\sum_{m=0}^{\infty} \Delta^{-m} \sum_{F \in \mathcal{F}_m(X,Y)} w(F) \;\le\; \sum_{i=0}^{\infty} \Delta^{-i} \sum_{F_1 \in \mathcal{F}_i(X \setminus x, Y)} w(F_1) \;\le\; 1 \qquad (5.8)
$$

by the inductive hypothesis. □

Examples 4.1 and 4.2 show that Proposition 5.1 is best possible (at least for a general set Y), even when $|X| = 1$.

Proof of Proposition 5.2. As before, we assume that $X \cap Y = \varnothing$ and we use induction on $k = |X|$. If $k = 0$ the result is trivial. Suppose next that $k = 1$ and $X = \{x\}$. Since $f_m(\{x\},\{y\}) = w_m^{\rm SAW}(x,y)$, we have $\sum\limits_{m=0}^{\infty} \Lambda^{-m} f_m(\{x\},\{y\}) \;\le\; 1$ for each $y \in Y$ by Proposition 4.3. Thus $\sum\limits_{m=0}^{\infty} \Lambda^{-m} f_m(\{x\}, Y) \;\le\; |Y|$ and the proposition holds for $k = 1$.

Suppose now that $k \ge 2$. Let x be any vertex in X; by assumption $x \notin Y$. By Lemma 5.5, given any $F \in \mathcal{F}_m(X, Y)$, we can decompose F into a forest $F_1 \in \mathcal{F}_i(X \setminus x, Y)$ and a path $P \in \mathcal{W}_{m-i}^{\mathrm{FPSAW}}(x, V(F_1))$; and each such pair (F_1, P) arises from a unique F. Since $w(F) = w(F_1)w(P)$, we have

$$\begin{aligned}
&\sum_{m=0}^{\infty} (m+|Y|)^{-(k-1)} \Lambda^{-m} \sum_{F \in \mathcal{F}_m(X,Y)} w(F) \\
&\quad \le \sum_{m=0}^{\infty} (m+|Y|)^{-(k-1)} \sum_{i=0}^{m} \Lambda^{-i} \sum_{F_1 \in \mathcal{F}_i(X \setminus x, Y)} w(F_1)\, \Lambda^{-(m-i)} \sum_{P \in \mathcal{W}_{m-i}^{\mathrm{FPSAW}}(x, V(F_1))} w(P) \\
&\quad = \sum_{i=0}^{\infty} \Lambda^{-i} \sum_{F_1 \in \mathcal{F}_i(X \setminus x, Y)} w(F_1) \sum_{j=0}^{\infty} \Lambda^{-j} (i+j+|Y|)^{-(k-1)} \sum_{P \in \mathcal{W}_{j}^{\mathrm{FPSAW}}(x, V(F_1))} w(P)\,.
\end{aligned} \tag{5.9}$$

Now, for each fixed F_1 we have $\sum_{j=0}^{\infty} \Lambda^{-j} \sum_{P \in \mathcal{W}_j^{\mathrm{FPSAW}}(x, V(F_1))} w(P) \le |V(F_1)|$ by the base case $k = 1$. Since $|V(F_1)| = i + |Y| \le i + j + |Y|$, we have

$$\begin{aligned}
\sum_{m=0}^{\infty} (m+|Y|)^{-(k-1)} \Lambda^{-m} \sum_{F \in \mathcal{F}_m(X,Y)} w(F) \;&\le\; \sum_{i=0}^{\infty} (i+|Y|)^{-(k-2)} \Lambda^{-i} \sum_{F_1 \in \mathcal{F}_i(X \setminus x, Y)} w(F_1) \\
&\le\; 1
\end{aligned} \tag{5.10}$$

by the inductive hypothesis. $\square$

When $|X| = 1$, Proposition 5.2 is in some sense best possible. To see this, take G to be a star, X to be the central vertex and Y to be the end-vertices: this gives $\Lambda = 1$, $f_1(X, Y) = |Y|$ and $f_m(X, Y) = 0$ for $m \ne 1$, so that $\sum_{m=0}^{\infty} \Lambda^{-m} f_m(X, Y) = |Y|$. When $|X| = k \ge 2$, by contrast, Proposition 5.2 is perhaps not best possible. If we take G to be the disjoint union of k isomorphic stars (again with central vertices in X and end-vertices in Y), we have $\sum_{m=0}^{\infty} \Lambda^{-m} f_m(X, Y) = (|Y|/k)^k$. This shows that if there is a universal upper bound on $\sum_{m=0}^{\infty} \Lambda^{-m} f_m(X, Y)$, the right-hand side has to grow at least like $(|Y|/|X|)^{|X|}$. We suspect the following conjectures are true:

Conjecture 5.6 *For all $X, Y \subseteq V(G)$ with $Y \ne \varnothing$, we have*

$$\sum_{m=0}^{\infty} \Lambda^{-m} f_m(X, Y) \;\le\; |Y|^{|X|}\,. \tag{5.11}$$

Conjecture 5.7 (a special case of Conjecture 5.6) *For all nonempty $X \subseteq V(G)$, we have*

$$\sum_{m=0}^{\infty} \Lambda^{-m} t_m(X) \;\le\; 1 \; . \tag{5.12}$$

5.2 The class $\mathcal{H}_m(X)$

We conclude this section by discussing a larger class of forests. For $X \subseteq V(G)$, let $\mathcal{H}_m(X)$ be the set of all m-edge forests F in G such that $L(F) \subseteq X \subseteq V(F)$. For integers $p, r \ge 1$, let $\mathcal{H}_m(X,p)$ be the set of all m-edge forests F in $\mathcal{H}_m(X)$ such that each component of F contains at least p vertices of X, and let $\mathcal{H}_m(X,p,r)$ be the set of all forests F in $\mathcal{H}_m(X,p)$ such that F has precisely r components. Put

$$h_m(X) \;=\; \sum_{F \in \mathcal{H}_m(X)} w(F) \; , \tag{5.13}$$

and define $h_m(X,p)$ and $h_m(X,p,r)$ similarly. Our next result uses Proposition 5.1 to bound $h_m(X,p,r)$ in terms of Δ:

Proposition 5.8 *Let $X \subseteq V(G)$ where $|X| = k \ge rp$. Then*

$$\sum_{m=0}^{\infty} \Delta^{-m} h_m(X,p,r) \;\le\; p^{-(r-1)} (k - rp + p)^{-1} \binom{k}{r} \; . \tag{5.14}$$

Proof. Choose $F \in \mathcal{H}_m(X,p,r)$. Let $\{X_1, X_2, \ldots, X_r\}$ be the partition of X determined by the components of F. Then $F \in \mathcal{F}_m(X,Y)$ for all sets Y such that $|Y \cap X_j| = 1$ for all $1 \le j \le r$. Hence there are precisely $\prod_{j=1}^{r} |X_j|$ different sets $Y \subseteq X$ such that $F \in \mathcal{F}_m(X,Y)$. Since $|X_j| \ge p$ for all j ($1 \le j \le r$) and $\sum_{j=1}^{r} |X_j| = k$, it follows that $\prod_{j=1}^{r} |X_j| \ge p^{r-1}(k - rp + p)$. Thus

$$\sum_{F \in \mathcal{H}_m(X,p,r)} w(F) \;\le\; p^{-(r-1)} (k - rp + p)^{-1} \sum_{\substack{Y \subseteq X \\ |Y| = r}} \; \sum_{F \in \mathcal{F}_m(X,Y)} w(F). \tag{5.15}$$

Using Proposition 5.1 we deduce

$$\begin{aligned} \sum_{m=0}^{\infty} \sum_{F \in \mathcal{H}_m(X,p,r)} \Delta^{-m} w(F) \;&\le\; p^{-(r-1)} (k - rp + p)^{-1} \sum_{m=0}^{\infty} \sum_{\substack{Y \subseteq X \\ |Y| = r}} \; \sum_{F \in \mathcal{F}_m(X,Y)} w(F) \\ &\le\; p^{-(r-1)} (k - rp + p)^{-1} \binom{k}{r} \; . \end{aligned} \tag{5.16}$$

□

Summing over r, we obtain:

Corollary 5.9 *Let $X \subseteq V(G)$ where $|X| = k \geq 1$. Then*

$$\sum_{m=0}^{\infty} \Delta^{-m} h_m(X,p) \;\leq\; \sum_{r=1}^{\lfloor k/p \rfloor} p^{-(r-1)} (k-rp+p)^{-1} \binom{k}{r} \qquad (5.17a)$$

$$\leq\; \left(1+\frac{1}{p}\right)^k - 1\,. \qquad (5.17b)$$

Here the crude upper bound (5.17b) is obtained from (5.17a) by replacing $(k-rp+p)^{-1}$ by p^{-1}. The true large-k asymptotic behavior of (5.17a) is $(1+1/p)^k k^{-1} p(p+1)[1+O(1/k)]$.

Taking $p=1$ in (5.17a) gives:

Corollary 5.10 *Let $X \subseteq V(G)$ where $|X| = k \geq 1$. Then*

$$\sum_{m=0}^{\infty} \Delta^{-m} h_m(X) \;\leq\; \frac{2(2^k-1)}{k+1}\,. \qquad (5.18)$$

Using a similar proof technique to that of Proposition 5.8, but using Proposition 5.2 instead of Proposition 5.1, we may deduce

Proposition 5.11 *Let $X \subseteq V(G)$ where $|X| = k \geq rp$. Then*

$$\sum_{m=0}^{\infty} (m+r)^{-(k-1)} \Lambda^{-m} h_m(X,p,r) \;\leq\; r p^{-(r-1)} (k-rp+p)^{-1} \binom{k}{r}\,. \qquad (5.19)$$

6 Counting Connected Subgraphs

For $X \subseteq V(G)$, let $\mathcal{C}_m(X)$ be the set of all m-edge subgraphs H in G (connected or not) such that

(C1) $X \subseteq V(H)$, and

(C2) each component of H contains at least one vertex in X.

Note in particular the following special cases:

- For any X, $\mathcal{C}_0(X)$ has as its single element the edgeless graph with vertex set X.
- $\mathcal{C}_0(\varnothing)$ has as its single element the empty graph, and $\mathcal{C}_m(\varnothing) = \varnothing$ for $m \geq 1$.
- $\mathcal{C}_m(\{x\})$ consists of the connected m-edge subgraphs that contain x.

Recall the definition $w(H) = \prod_{e \in E(H)} w_e$. Define the weighted counts

$$c_m(X) \;=\; \sum_{H \in \mathcal{C}_m(X)} w(H) \;. \tag{6.1}$$

We then have the following bound in terms of maximum weighted degree, which is a slight extension of [12, Proposition 4.5]:

Proposition 6.1 *Whenever $|X| = k \geq 0$ we have*

$$c_m(X) \;\leq\; C(m,k)\,\Delta^m \tag{6.2}$$

where

$$C(m,k) \;=\; \begin{cases} k(m+k)^{m-1}/m! & \textit{for } k \neq 0 \\ \delta_{m0} & \textit{for } k = 0 \end{cases} \tag{6.3}$$

Before beginning the proof of Proposition 6.1, let us note some facts about the numbers $C(m,k)$:

(a) For each integer $m \geq 0$, $C(m,k)$ is a polynomial of degree m in k.

(b) For each integer $m \geq 0$, $C(m,k)$ is an increasing function of k for $k \geq 0$.

(c) Generating function: If $\mathsf{C}(z)$ solves the equation

$$\mathsf{C}(z) \;=\; e^{z\mathsf{C}(z)} \;, \tag{6.4}$$

then

$$\mathsf{C}(z)^k \;=\; \sum_{m=0}^{\infty} C(m,k)\, z^m \tag{6.5}$$

for all k (integer or not); this follows from the Lagrange inversion formula. Moreover, the series (6.5) is absolutely convergent for $|z| \leq 1/e$ and satisfies $\mathsf{C}(1/e) = e$.

(d) For integer $k \geq 1$,

$$C(m,k) \;=\; \sum_{\substack{m_1, \ldots, m_k \geq 0 \\ m_1 + \cdots + m_k = m}} \prod_{i=1}^{k} C(m_i, 1) \;. \tag{6.6}$$

This is an immediate consequence of (6.5).

(e) For all k and z (integer or not),

$$C(m,k) \;=\; \sum_{f=0}^{m} \frac{z^f}{f!}\, C(m-f, k-z+f)\;. \tag{6.7}$$

This can easily be verified by direct calculation using the binomial formula:

$$\begin{aligned}
&\sum_{f=0}^{m} \frac{z^f}{f!}\, \frac{(k-z+f)(m+k-z)^{m-f-1}}{(m-f)!} \\
&\qquad = \; \frac{1}{m!} \sum_{f=0}^{m} \binom{m}{f} (k-z+f) z^f (m+k-z)^{m-f-1} \\
&\qquad = \; \frac{1}{m!} \left(1 - \frac{d}{dk}\right) \sum_{f=0}^{m} \binom{m}{f} z^f (m+k-z)^{m-f} \\
&\qquad = \; \frac{1}{m!} \left(1 - \frac{d}{dk}\right) (m+k)^m \\
&\qquad = \; \frac{1}{m!} \left[(m+k)^m \;-\; m(m+k)^{m-1}\right] \\
&\qquad = \; \frac{k(m+k)^{m-1}}{m!}\;.
\end{aligned} \tag{6.8}$$

(f) For each fixed $k > 0$, we have

$$C(m,k) \;=\; e^m m^{-3/2} \frac{k e^k}{\sqrt{2\pi}} \left[1 + O(1/m)\right] \tag{6.9}$$

as $m \to \infty$. This is an immediate consequence of Stirling's formula.

In proving Proposition 6.1, we will use the following two facts about the weighted counts $c_m(X)$ for $X = \{x_1, x_2, \ldots, x_k\} \subseteq V(G)$ where $x_1, x_2, \ldots, x_k$ are all distinct:

Fact 1. $c_m(X) \le \displaystyle\sum_{\substack{m_1, \ldots, m_k \ge 0 \\ m_1 + \cdots + m_k = m}} \prod_{i=1}^{k} c_{m_i}(x_i)\;.$

Proof. Trivial when $k = 1$. For $k \ge 2$, construct a weight-preserving bijection $\mathbf{F}$ of $\mathcal{C}_m(X)$ onto a subset of $\displaystyle\bigcup_{\substack{m_1, \ldots, m_k \ge 0 \\ m_1 + \cdots + m_k = m}} \prod_{i=1}^{k} \mathcal{C}_{m_i}(x_i)$ as follows: Given $H \in \mathcal{C}_m(X)$, define $\mathbf{F}(H)_i$ to be the component of H containing x_i if this component contains no vertex $x_{i'}$ with $i' < i$, and the graph with vertex set $\{x_i\}$ and no edges, otherwise. $\square$

Fact 2. $c_m(X) \le \sum\limits_{F \subseteq C(x_1)} w(F)\, c_{m-|F|}((X - x_1) \cup Y^F)$ where $C(x_1)$ is shorthand for the cocycle $E(\{x_1\}, \{x_1\}^c)$, and Y^F denotes the set of endpoints other than x_1 of the edges in F. [Of course, the sum can be limited to sets F having $|F| \le m$, since $c_m(X) = 0$ for $m < 0$.]

Proof. We classify the subgraphs $H \in \mathcal{C}_m(X)$ according to $F \equiv E(H) \cap C(x_1)$. So, for each $F \subseteq C(x_1)$, let $\mathcal{C}_m(X; F)$ be the set of all $H \in \mathcal{C}_m(X)$ that have $E(H) \cap C(x_1) = F$. For $H \in \mathcal{C}_m(X; F)$, we let $H' = H \setminus x_1$. Then $|E(H')| = m - |F|$ and $H' \in \mathcal{C}_{m-|F|}((X - x_1) \cup Y^F)$. This gives an injective map from $\mathcal{C}_m(X; F)$ into $\mathcal{C}_{m-|F|}((X - x_1) \cup Y^F)$. Fact 2 now follows since $w(H) = w(F)w(H')$. $\square$

We now give two alternative proofs of Proposition 6.1. The first proof uses Fact 2 for $k = 1$ only, together with Fact 1; it leads to a *nonlinear* recursion whose solution is $C(m, 1)$, namely (6.7) with $k = z = 1$ combined with (6.6). The second proof uses Fact 2 for all k, but does not use Fact 1; it leads to a *linear* recursion whose solution is $C(m, k)$, namely (6.7) with $z = 1$.

We will need the following elementary result:

Lemma 6.2 *Let S be a set in which each element $e \in S$ is given a nonnegative real weight w_e. Then, for each integer $f \ge 0$, we have*

$$\sum_{\substack{F \subseteq S \\ |F| = f}} \prod_{e \in F} w_e \;\le\; \frac{1}{f!} \left(\sum_{e \in S} w_e \right)^{f} . \tag{6.10}$$

First proof of Proposition 6.1. Consider first the case $k = 1$, hence $X = \{x\}$. We use induction on m. The proposition holds trivially when $m = 0$, so let us assume that $m \ge 1$. We apply Fact 2 with $k = 1$, and observe that the term $F = \varnothing$ contributes zero when $m \ge 1$ [since $c_{m-|F|}(Y^F) = c_m(\varnothing) = 0$]; we therefore have

$$c_m(x) \;\le \sum_{\substack{\varnothing \ne F \subseteq C(x) \\ |F| \le m}} w(F)\, c_{m-|F|}(Y^F) \,. \tag{6.11}$$

Applying Fact 1 to bound $c_{m-|F|}(Y^F)$, we obtain

$$c_m(x) \;\le \sum_{\substack{\varnothing \ne F \subseteq C(x) \\ |F| \le m}} w(F) \sum_{\substack{m_1, \ldots, m_{|Y^F|} \ge 0 \\ m_1 + \cdots + m_{|Y^F|} = m - |F|}} \prod_{i=1}^{|Y^F|} c_{m_i}(v_i) \tag{6.12}$$

where $Y^F = \{v_1, \ldots, v_{|Y^F|}\}$. Note that we have $m_i < m$ for all i since $|F| \geq 1$. We can therefore apply the inductive hypothesis $c_{m_i}(v_i) \leq C(m_i, 1)\, \Delta^{m_i}$ to obtain

$$\begin{aligned}
c_m(x) &\leq \sum_{\substack{\varnothing \neq F \subseteq C(x) \\ |F| \leq m}} w(F) \sum_{\substack{m_1, \ldots, m_{|Y^F|} \geq 0 \\ m_1 + \cdots + m_{|Y^F|} = m - |F|}} \prod_{i=1}^{|Y^F|} C(m_i, 1)\, \Delta^{m_i} \\
&= \sum_{\substack{\varnothing \neq F \subseteq C(x) \\ |F| \leq m}} w(F)\, C(m - |F|,\, |Y^F|)\, \Delta^{m-|F|} \\
&\leq \sum_{\substack{\varnothing \neq F \subseteq C(x) \\ |F| \leq m}} w(F)\, C(m - |F|,\, |F|)\, \Delta^{m-|F|} \qquad (6.13)
\end{aligned}$$

where the second line used the identity (6.6), and the last step used $|Y^F| \leq |F|$ and the fact that $C(m, k)$ is an increasing function of k.[3] Using Lemma 6.2, we have

$$\begin{aligned}
c_m(x) &\leq \sum_{f=1}^{m} \frac{\Delta^f}{f!}\, C(m - f, f)\, \Delta^{m-f} \\
&= \sum_{f=0}^{m} \frac{\Delta^f}{f!}\, C(m - f, f)\, \Delta^{m-f} \\
&= C(m, 1)\, \Delta^m\,, \qquad (6.14)
\end{aligned}$$

where the second line used $C(m, 0) = 0$ for $m \geq 1$, and the last line used identity (6.7) with $k = z = 1$. This proves Proposition 6.1 for $k = 1$.

The result for general k now follows using Fact 1 and the identity (6.6). □

SECOND PROOF OF PROPOSITION 6.1. We use induction on $m+k$. The proposition holds trivially when $m = 0$, so we assume $m \geq 1$. Apply Fact 2 for general k; because the right-hand side involves quantities $c_{m-|F|}((X - x_1) \cup Y^F)$ with $m - |F| + |(X - x_1) \cup Y^F| \leq m + k - 1$, we can apply the inductive hypothesis $c_{m-|F|}((X - x_1) \cup Y^F) \leq C(m - |F|, |(X - x_1) \cup Y^F|)\, \Delta^{m-|F|}$ to conclude that

$$\begin{aligned}
c_m(X) &\leq \sum_{F \subseteq C(x_1)} w(F)\, C(m - |F|,\, |(X - x_1) \cup Y^F|)\, \Delta^{m-|F|} \\
&\leq \sum_{F \subseteq C(x_1)} w(F)\, C(m - |F|,\, k - 1 + |F|)\, \Delta^{m-|F|}
\end{aligned}$$

---

[3] If G is a simple graph, we have $|Y^F| = |F|$. But if G has multiple edges, all we can say is that $|Y^F| \leq |F|$.

$$\begin{aligned} &\leq \sum_{f=0}^{m} \frac{\Delta^f}{f!}\, C(m-f,\, k-1+f)\, \Delta^{m-f} \\ &= C(m,k)\, \Delta^m\ , \end{aligned} \tag{6.15}$$

where the second line used $|(X - x_1) \cup Y^F| \leq k-1+|Y^F| \leq k-1+|F|$ and the fact that $C(m,k)$ is increasing with k, the third line used Lemma 6.2, and the last line used identity (6.7) with $z = 1$. $\square$

Let us now show that Proposition 6.1 is best possible:

Example 6.1. Let $\mathbf{T}_r$ be the infinite r-regular tree, let $x_1, \ldots, x_k \in V(\mathbf{T}_r)$ satisfy $\mathrm{dist}(x_i, x_j) > m$ for $i \neq j$, and let G be the subtree of $\mathbf{T}_r$ induced by the vertices lying at a distance at most M from the set $\{x_1, \ldots, x_k\}$, where M is any fixed number greater than or equal to m. Let all edge weights w_e equal Δ/r. Then by a slight generalization of the computation in [12, proof of Proposition 4.2], one shows that

$$c_m(x_1, \ldots, x_k) \;=\; \left(\frac{\Delta}{r}\right)^m \frac{kr}{m} \binom{kr + (r-1)m - 1}{m-1}\ , \tag{6.16}$$

which in the limit $r \to \infty$ with Δ fixed tends to $C(m,k)\, \Delta^m$. $\square$

Example 6.2. Let G be the complete graph K_n, and let all edge weights w_e equal $\Delta/(n-1)$. Consider first the case $k = 1$: let us count the subset of $\mathcal{C}_m(\{x\})$ consisting of the m-edge *trees* containing x. The number of such trees is $\binom{n-1}{m}(m+1)^{m-1}$, where the first factor counts the number of ways we can choose m additional vertices and the second factor counts the number of trees on $m+1$ labelled vertices. We therefore have

$$c_m(x) \;\geq\; \binom{n-1}{m}(m+1)^{m-1} \left(\frac{\Delta}{n-1}\right)^m\ , \tag{6.17}$$

which in the limit $n \to \infty$ with m fixed tends to $C(m,1)\, \Delta^m$.

Now consider general k: let $x_1, \ldots, x_k$ be distinct vertices, and let us count the subset of $\mathcal{C}_m(x_1, \ldots, x_k)$ consisting of the m-edge k-component *forests* in which each component contains exactly one x_i. By a similar counting argument we have

$$\begin{aligned} &c_m(x_1, \ldots, x_k) \;\geq \\ &\qquad \left(\frac{\Delta}{n-1}\right)^m \sum_{\substack{m_1, \ldots, m_k \geq 0 \\ m_1 + \cdots + m_k = m}} \prod_{i=1}^{k} \binom{n-k-m_1-\ldots-m_{i-1}}{m_i} (m_i+1)^{m_i - 1}\ , \end{aligned} \tag{6.18}$$

which in the limit $n \to \infty$ with m fixed tends to $C(m,k)\, \Delta^m$ [using (6.6)]. $\square$

Note the difference between the "pointwise" bounds of Propositions 4.1 and 6.1 and the "generating-function" bounds of Propositions 4.2, 4.3, 5.1 and 5.2. In the former cases, the bound can be saturated simultaneously for all (or arbitrarily many) m — as shown by the preceding two examples — so nothing can be gained by summing over m. In the latter cases, by contrast, the pointwise bound can be saturated only for one m at a time (cf. Example 4.3 after Corollary 4.5), so the generating-function bound is stronger. The fundamental difference between the two situations is that the subgraphs considered in Propositions 4.1 and 6.1 are "tied down only at one end" (and hence can grow freely in all directions), while those in Propositions 4.2, 4.3, 5.1 and 5.2 are "tied down at all the leaves".

7 Counting Blocks, Block Paths, Block Trees and Block Forests

We now seek an analogue of Proposition 6.1 using maxmaxflow in place of maximum degree. Unfortunately, no such bound is possible for the families $\mathcal{C}_m(X)$: a simple counterexample is provided by the stars $G = K_{1,r}$, which have maximum degree r but maxmaxflow 1 (independent of r); letting x be the central vertex of the star, we have $|\mathcal{C}_1(\{x\})| = r$, which is unbounded as $r \to \infty$. The same thing happens for wheels $G = K_1 + C_r$, so it is no help to assume that G is 2-connected.

To get a bound in terms of maxmaxflow, we need to restrict attention to a suitably chosen proper *subfamily* of $\mathcal{C}_m(X)$: roughly speaking, we need to count subgraphs H that possess suitable "2-connectedness properties". After much experimentation, we settled on the family $\mathcal{BT}_m(X)$ of "block trees" and the families $\mathcal{BF}_m(X,Y)$ and $\mathcal{BF}^*_m(X,Y)$ of "block forests", to be defined in Section 7.1. However, simpler but weaker results can be obtained using a different (and slightly larger) family $\mathcal{B}_m(X)$ of block forests, to be defined in Section 7.2. These two subsections are independent of each other and can be read in either order.

7.1 The classes $\mathcal{BT}_m(X)$, $\mathcal{BF}_m(X,Y)$ and $\mathcal{BF}^*_m(X,Y)$

Let H be a not-necessarily-connected graph. We recall that a vertex of H is called an *internal vertex* of H if it is not a cut vertex of H. Now let B be a block of H. We denote by $\mathrm{Int}(B,H)$ the set of internal vertices of H that belong to B. We say that B is an *isolated block* of H if it contains no cut vertices of H, and an *end block* of H if it contains exactly one cut vertex of H. Finally, if $x, y \in V(H)$ with $x \neq y$, we say that H is an *xy-block path* if H is connected and is either a block (hence an isolated block of itself) or else has exactly two end blocks B_1, B_2 with $x \in \mathrm{Int}(B_1,H)$ and $y \in \mathrm{Int}(B_2,H)$.

For any nonempty $X \subseteq V(G)$, let $\mathcal{BT}_m(X)$ be the set of all m-edge subgraphs H in G such that

(BT1) $X \subseteq V(H)$;

(BT2) H is connected;

(BT3) each end block B of H contains at least one vertex of X as an internal vertex [that is, $\mathrm{Int}(B,H) \cap X \neq \varnothing$]; and

(BT4) if H is a block, then either H is an isolated vertex [hence $V(H) = X = \{x\}$] or else H contains at least two vertices of X.

By analogy with the set $\mathcal{T}_m(X)$, which consists of trees whose leaves are "tied down" on the set X, we think of the elements of $\mathcal{BT}_m(X)$ as *block trees* whose end blocks are "tied down" on the set X. [In particular, we have $\mathcal{T}_m(X) \subseteq \mathcal{BT}_m(X)$.] Note the following special cases:

- $\mathcal{BT}_0(\{x\})$ has as its single element the edgeless graph with vertex set $\{x\}$, and $\mathcal{BT}_m(\{x\}) = \varnothing$ for $m \geq 1$.
- When $x \neq y$, $\mathcal{BT}_m(\{x,y\})$ is the set of m-edge xy-block paths.
- $\mathcal{BT}_0(X) = \varnothing$ for $|X| \geq 2$.

For $X, Y \subseteq V(G)$ with $Y \neq \varnothing$, let $\mathcal{BF}_m(X,Y)$ be the set of all m-edge subgraphs H in G (connected or not) such that

(BF1) $X \cup Y \subseteq V(H)$;

(BF2) each component of H contains exactly one vertex of Y;

(BF3) each end block B of H contains at least one vertex of $X \cup Y$ as an internal vertex [that is, $\mathrm{Int}(B,H) \cap (X \cup Y) \neq \varnothing$]; and

(BF4) each isolated block of H is either an isolated vertex belonging to Y or else contains at least two vertices of $X \cup Y$.

[Note that $\mathcal{BF}_m(X,Y) = \mathcal{BF}_m(X \setminus Y, Y)$, so we can assume without loss of generality, if desired, that $X \cap Y = \varnothing$.] By analogy with the set $\mathcal{F}_m(X,Y)$, which consists of forests whose leaves are "tied down" on the set X and whose components are "tied down" on single elements of the set Y, we think of the elements of $\mathcal{BF}_m(X)$ as *block forests* whose end blocks are "tied down" on the set X and whose components are "tied down" on single elements of the set Y. [In particular, we have $\mathcal{F}_m(X,Y) \subseteq \mathcal{BF}_m(X,Y) \subseteq \mathcal{C}_m(Y)$.] Note the following special cases:

- If $m = 0$, then $\mathcal{BF}_0(X,Y) = \varnothing$ whenever $X \not\subseteq Y$, and $\mathcal{BF}_0(X,Y)$ has as its single element the edgeless graph with vertex set Y whenever $X \subseteq Y$.
- If $X = \varnothing$ (or more generally if $X \subseteq Y$), then $\mathcal{BF}_0(X,Y)$ has as its single element the edgeless graph with vertex set Y, and $\mathcal{BF}_m(X,Y) = \varnothing$ for $m \geq 1$.

- If $X = \{x\}$ with $x \notin Y$, then each $H \in \mathcal{BF}_m(\{x\}, Y)$ is the disjoint union of an m-edge xy-block path for some $y \in Y$ (this component avoiding the set $Y \setminus y$) and the collection $Y \setminus y$ of isolated vertices.
- For $Y = \{y\}$, we have $\mathcal{BF}_m(X, \{y\}) = \mathcal{BT}_m(X \cup \{y\})$.

Recall that $w(H) = \prod_{e \in E(H)} w_e$. Define the weighted counts

$$bt_m(X) \;=\; \sum_{H \in \mathcal{BT}_m(X)} w(H) \tag{7.1}$$

$$bf_m(X, Y) \;=\; \sum_{H \in \mathcal{BF}_m(X,Y)} w(H) \tag{7.2}$$

We will obtain the following bounds on $bf_m(X, Y)$ in terms of Δ and Λ:

Proposition 7.1 *For all $X, Y \subseteq V(G)$ with $Y \neq \varnothing$, we have*

$$\sum_{m=0}^{\infty} \left(\frac{\Delta}{\ln 2} \right)^{-m} bf_m(X, Y) \;\le\; 1\,. \tag{7.3}$$

Proposition 7.2 *For all $X, Y \subseteq V(G)$ with $Y \neq \varnothing$ and all $\alpha \in (1, 2]$, we have*

$$\sum_{m=0}^{\infty} \left(\frac{\alpha\Lambda}{\ln \alpha} \right)^{-m} bf_m(X, Y) \;\le\; \alpha^{|Y|-1}\,. \tag{7.4}$$

Using the identity $\mathcal{BF}_m(X, \{y\}) = \mathcal{BT}_m(X \cup \{y\})$, we immediately obtain the following corollaries for block trees:

Corollary 7.3 *For all nonempty $X \subseteq V(G)$, we have*

$$\sum_{m=0}^{\infty} \left(\frac{\Delta}{\ln 2} \right)^{-m} bt_m(X) \;\le\; 1\,. \tag{7.5}$$

Corollary 7.4 *For all nonempty $X \subseteq V(G)$, we have*

$$\sum_{m=0}^{\infty} \left(\frac{2\Lambda}{\ln 2} \right)^{-m} bt_m(X) \;\le\; 1\,. \tag{7.6}$$

Corollary 7.3 could be proved directly using the same proof technique as for Proposition 7.1. It is curious to note, however, that we have been unable to find a direct proof of Corollary 7.4; our proof of Proposition 7.2 employs an inner induction on $|Y|$, and thus inevitably passes through disconnected graphs.

As an immediate consequence of Corollary 7.4, we obtain:

Corollary 7.5 *Fix a weighted graph* $(G, \mathbf{w})$ *and an edge* $e \in E(G)$*. Let* $b_m(e)$ *be the sum of the weights of the 2-connected* m*-edge subgraphs of* G *containing* e*. Then*

$$\sum_{m=0}^{\infty} \left(\frac{2\Lambda(G-e, \mathbf{w})}{\ln 2} \right)^{-m} b_m(e) \;\le\; 1 \; . \tag{7.7}$$

Proof of Corollary 7.5, assuming Corollary 7.4. If $e = xy$, this follows immediately from Corollary 7.4 applied to $G - e$ by taking $X = \{x, y\}$. $\square$

Our proof of Proposition 7.1 combines ideas from the proofs of Propositions 5.1 and 6.1: we use an inner induction on $|X|$ as a substitute for the "point-to-set" bound of Proposition 4.2, and we use a "cutting" argument similar to that employed in the proofs of Propositions 4.2 and 6.1 to handle the case $|X| = 1$. In the inductive step we shall use the following analogue of Lemma 5.5 to split a block forest with k end blocks into a block forest with $k-1$ end blocks and a block path:

Lemma 7.6 *Let* G *be a graph, let* $X, Y \subseteq V(G)$ *with* $Y \neq \varnothing$*, let* $x \in X \setminus Y$*, and let* $H \in \mathcal{BF}_m(X, Y)$*. Let* H_1 *be the convex hull of* $(X \setminus x) \cup Y$ *in* H*, and let* H_2 *be the convex hull of* $\{x\} \cup V(H_1)$ *in* $H \setminus E(H_1)$*. Then* H *is the edge-disjoint union of* H_1 *and* H_2*; and for some* i $(0 \le i \le m)$*, we have* $H_1 \in \mathcal{BF}_i(X \setminus x, Y)$ *and* $H_2 \in \mathcal{BF}_{m-i}(\{x\}, V(H_1))$*. Moreover, the map* $H \mapsto (H_1, H_2)$ *is an injection.*

Note the slight change of perspective from Lemma 5.5: here H_2 is not an xy-block path for some $y \in V(H_1)$, but rather the union of such a block path with the collection $V(H_1) \setminus y$ of isolated vertices. In particular, we have $V(H_1) \subseteq V(H_2)$. However, modulo this change, this decomposition reduces to that of Lemma 5.5 in the special case where $H \in \mathcal{F}_m(X, Y)$.

Proof of Proposition 7.1. As noted above, we can assume without loss of generality that $X \cap Y = \varnothing$. Let $c = 1/\ln 2$. We shall show that

$$\sum_{m=0}^{M} (c\Delta)^{-m}\, bf_m(X, Y) \;\le\; 1 \tag{7.8}$$

for all $M \ge 0$, by using an outer induction on M and an inner induction on $|X|$. The base case $M = 0$ and $|X|$ arbitrary holds by the first remark after the definition of $\mathcal{BF}_m(X, Y)$. The case when $X = \varnothing$ and M is arbitrary holds by the second remark following the definition of $\mathcal{BF}_m(X, Y)$. Hence we may suppose that $M \ge 1$ and $|X| \ge 1$. Our inductive argument consists of two steps:

Step 1. Proof that if (7.8) holds for all $|X|$ and all M' with $0 \le M' < M$, then it also holds for $|X| = 1$ and M. This step uses a "cutting" argument.

Step 2. Proof that if (7.8) holds for all $|X'|$ with $1 \le |X'| < |X|$ and some given M, then it holds for $|X|$ and the same M. This step uses Lemma 7.6.

Step 1. Suppose that (7.8) holds for all $|X|$ and all M' with $0 \le M' < M$. Now let $X = \{x\}$ (note that $x \notin Y$ by assumption) and consider a subgraph $H \in \mathcal{BF}_m(\{x\}, Y)$ for some m. Note that H is the disjoint union of an xy-block path H' for some $y \in Y$ and the collection $Y \setminus y$ of isolated vertices. In particular, x is neither an isolated vertex nor a cut vertex of H. Let F be the set of edges of H incident with x, let $f = |F|$ (≥ 1 because x is not isolated), and let U^F be the set of end-vertices of edges in F distinct from x. Then H' remains connected under deletion of x (because x is not a cut vertex).

We shall show that $H \setminus x \in \mathcal{BF}_{m-f}(U^F, Y)$. For each end block B of $H' \setminus x$, we have $\mathrm{Int}(B, H' \setminus x) \cap (U_F \cup \{y\}) \neq \varnothing$, otherwise B would be an end block of H with $\mathrm{Int}(B, H) \cap (\{x\} \cup Y) \neq \varnothing$. Furthermore, if $H' \setminus x$ is non-separable, then either $V(H' \setminus x) = \{y\}$, or $H' \setminus x$ contains at least two vertices of $U_F \cup \{y\}$); otherwise $H' \setminus x$ would itself be an end block of H with $\mathrm{Int}(H' \setminus x, H) \cap (\{x\} \cup Y) \neq \varnothing$. Thus $H \setminus x \in \mathcal{BF}_{m-f}(U^F, Y)$.

Let C be the set of edges of G incident to x. For each nonempty $F \subseteq C$, let $\mathcal{BF}_m(\{x\}, Y; F)$ be the set of all subgraphs $H \in \mathcal{BF}_m(\{x\}, Y)$ such that $E(H) \cap C = F$. Then the map $H \mapsto H \setminus x$ is an injection from $\mathcal{BF}_m(\{x\}, Y; F)$ into $\mathcal{BF}_{m-f}(U^F, Y)$, and $w(H) = w(F) w(H \setminus x)$. Thus

$$bf_m(\{x\}, Y) \;\le\; \sum_{\varnothing \neq F \subseteq C} w(F)\, bf_{m-f}(U^F, Y)\ . \tag{7.9}$$

Hence

$$\begin{aligned}
\sum_{m=0}^{M} (c\Delta)^{-m}\, bf_m(\{x\}, Y) \;&=\; \sum_{m=0}^{M} (c\Delta)^{-m} \sum_{\varnothing \neq F \subseteq C} w(F)\, bf_{m-f}(U^F, Y) \\
&=\; \sum_{\varnothing \neq F \subseteq C} w(F)\, (c\Delta)^{-f} \sum_{j=0}^{M-f} (c\Delta)^{-j}\, bf_j(U^F, Y) \\
&\le\; \sum_{\varnothing \neq F \subseteq C} w(F)\, (c\Delta)^{-f} \qquad\qquad (7.10)
\end{aligned}$$

by the outer inductive hypothesis on M. Using Lemma 6.2, we deduce that

$$\sum_{m=0}^{M} (c\Delta)^{-m}\, bf_m(\{x\}, Y) \;\le\; \sum_{f=1}^{\infty} (c\Delta)^{-f} \sum_{\substack{F \subseteq C \\ |F| = f}} w(F)$$

$$
\begin{aligned}
&\leq \sum_{f=1}^{\infty} (c\Delta)^{-f} \frac{\Delta^f}{f!} \\
&= e^{1/c} - 1 \;=\; 1
\end{aligned}
$$

since $c = 1/\ln 2$. This proves Step 1.

Step 2. Suppose that (7.8) holds for all $|X'|$ with $1 \leq |X'| < |X|$ and some given M. Then $|X| \geq 2$. Choose arbitrarily some $x \in X$ (recall again that $x \notin Y$). Given $H \in \mathcal{BF}_m(X, Y)$, by Lemma 7.6 we may decompose H into $H_1 \cup H_2$ where $H_1 \in \mathcal{BF}_i(X \backslash x, Y)$ and $H_2 \in \mathcal{BF}_{m-i}(\{x\}, V(H_1))$. Applying the inductive hypothesis that (7.8) holds both for $X' = X \setminus x$ and for $X' = \{x\}$ with the given M, we may deduce that

$$
\begin{aligned}
&\sum_{m=0}^{M} (c\Delta)^{-m} bf_m(X, Y) \\
&\quad= \sum_{m=0}^{M} (c\Delta)^{-m} \sum_{H \in \mathcal{BF}_m(X,Y)} w(H) \\
&\quad\leq \sum_{m=0}^{M} \sum_{i=0}^{m} (c\Delta)^{-i} \sum_{H_1 \in \mathcal{BF}_i(X\backslash x, Y)} w(H_1)\, (c\Delta)^{-(m-i)} \sum_{H_2 \in \mathcal{BF}_{m-i}(\{x\}, V(H_1))} w(H_2) \\
&\quad\leq \sum_{i=0}^{M} (c\Delta)^{-i} \sum_{H_1 \in \mathcal{BF}_i(X\backslash x, Y)} w(H_1) \sum_{j=0}^{M-i} (c\Delta)^{-j} \sum_{H_2 \in \mathcal{BF}_j(\{x\}, V(H_1))} w(H_2) \\
&\quad\leq 1\,. \qquad (7.11)
\end{aligned}
$$

This completes the proof of Step 2 and hence of the proposition. □

Remarks. 1. Step 2 works for any value of c. The specific value $c = 1/\ln 2$ enters only in Step 1.

2. Step 2 is "almost" unnecessary; we can "almost" apply Step 1 for any X. The trouble is that if $|X| > 1$ we might find that x is a cut vertex of H. In this case $H \setminus x \notin \mathcal{BF}_{m-f}(U^F, Y)$ since it will have one or more components containing no vertices of Y.

Example 7.1. Let G be the graph $K_2^{(s)}$ consisting of a pair of vertices x, y joined by s parallel edges. Set all edge weights $w_e = \Delta/s$. Let us consider $bt_m(\{x, y\}) = bf_m(\{x\}, \{y\})$. The generating function is

$$
\sum_{m=0}^{\infty} \zeta^m bf_m(\{x\}, \{y\}) \;=\; \left(1 + \frac{\Delta}{s}\zeta\right)^s \;-\; 1\,, \qquad (7.12)
$$

which is an increasing function of s (at fixed Δ and ζ) and tends to $e^{\Delta\zeta}-1$ as $s \to \infty$. It follows that Proposition 7.1 and Corollary 7.3 are sharp in the sense that $(\ln 2)/\Delta$ is the maximal value of ζ that allows an upper bound of 1. $\square$

Example 7.2. Let G be the disjoint union of n copies of $K_2^{(s)}$; let X contain one vertex from each copy, and let Y be the remaining vertices. Set all edge weights $w_e = \Delta/s$. Then

$$\sum_{m=0}^{\infty} \zeta^m bf_m(X,Y) \;=\; \left[\left(1+\frac{\Delta}{s}\zeta\right)^s \,-\, 1\right]^n \;\overset{s\to\infty}{\longrightarrow}\; (e^{\Delta\zeta}-1)^n \,. \tag{7.13}$$

So $(\ln 2)/\Delta$ is the maximal value of ζ that allows *any* finite upper bound that is independent of $|X|$ and $|Y|$. $\square$

Might it be possible to bound $\sum_{m=0}^{\infty} \zeta^m bf_m(X,Y)$ for some $\zeta > (\ln 2)/\Delta$ if we allow the right-hand side to depend on $|X|$ and $|Y|$? We doubt it; but all we can say for sure, at present, is that ζ cannot exceed $(2\ln 2)/\Delta$:

Example 7.3. Let G be the graph $P_n^{(s)}$ obtained from the n-edge path ($n \ge 2$) by replacing each edge by s parallel edges. Set all edge weights $w_e = \Delta/(2s)$. Let $x, y \in V(P_n^{(s)})$ with $\operatorname{dist}(x,y) = \ell$. Then

$$\sum_{m=0}^{\infty} \zeta^m bf_m(\{x\},\{y\}) \;=\; \left[\left(1+\frac{\Delta}{2s}\zeta\right)^s \,-\, 1\right]^\ell \;\overset{s\to\infty}{\longrightarrow}\; (e^{\Delta\zeta/2}-1)^\ell \,. \tag{7.14}$$

Since ℓ can be arbitrarily large, a universal upper bound on $\sum_{m=0}^{\infty} \zeta^m bf_m(X,Y)$ is impossible for $\zeta > (2\ln 2)/\Delta$, even when $|X| = |Y| = 1$. $\square$

We have been unable to obtain a bound for $bf_m(X,Y)$ in terms of Λ (i.e., Proposition 7.2 or something like it) by extending the proof technique of Proposition 5.2 in a similar way to the above proof. The problem is that a universal "point-to-set" bound of the form $\sum_{m=0}^{\infty} (c\Lambda)^{-m} bf_m(X,Y) \le 1$ (with a right-hand side that is independent of $|Y|$) is simply not valid for any constant c: it suffices to consider $G = K_{1,r}$ with $r > c$. If, on the other hand, we try to adapt the proof technique of Proposition 7.1 by using an inductive hypothesis of the form $\sum_{m=0}^{\infty} (m+|Y|)^{-(|X|-1)} (c\Lambda)^{-m} bf_m(X,Y) \le |Y|$ (similar to that of Proposition 5.2), then we are unable to carry through Step 1 because of the increase in the size of X when $\{x\}$ is replaced by U^F. Instead, our proof of Proposition 7.2 will rely solely on a "cutting" argument rather than using Lemma 7.6. In order for the induction to go through, we need to work with a slightly larger family of graphs than $\mathcal{BF}_m(X,Y)$.

For $X, Y \subseteq V(G)$ with $Y \neq \varnothing$, let $\mathcal{BF}_m^*(X,Y)$ be the set of all m-edge subgraphs H in G (connected or not) such that

(BF1) $X \cup Y \subseteq V(H)$;

(BF2*) each component of H contains at least one vertex of Y;

(BF3) each end block B of H contains at least one element of $X \cup Y$ as an internal vertex [that is, $\mathrm{Int}(B, H) \cap (X \cup Y) \neq \varnothing$]; and

(BF4) each isolated block of H is either an isolated vertex belonging to Y or else contains at least two vertices of $X \cup Y$.

The only change from $\mathcal{BF}_m(X, Y)$ is, therefore, that each component of H must contain *at least one* vertex of Y, rather than exactly one. We have $\mathcal{BF}_m(X, Y) \subseteq \mathcal{BF}^*_m(X, Y) \subseteq \mathcal{C}_m(Y)$. Since $\mathcal{BF}^*_m(X, Y) = \mathcal{BF}^*_m(X \setminus Y, Y)$, we can assume without loss of generality, if desired, that $X \cap Y = \varnothing$. Note the special cases:

- If $m = 0$, then $\mathcal{BF}^*_0(X, Y) = \varnothing$ whenever $X \not\subseteq Y$, and $\mathcal{BF}^*_0(X, Y)$ has as its single element the edgeless graph with vertex set Y whenever $X \subseteq Y$.
- If $Y = \{y\}$, then $\mathcal{BF}^*_m(X, \{y\}) = \mathcal{BF}_m(X, \{y\}) = \mathcal{BT}_m(X \cup \{y\})$.

Define the weighted counts

$$bf^*_m(X, Y) \;=\; \sum_{H \in \mathcal{BF}^*_m(X,Y)} w(H)\; . \tag{7.15}$$

Since $\mathcal{BF}_m(X, Y) \subseteq \mathcal{BF}^*_m(X, Y)$, Proposition 7.2 will follow from the stronger result:

Proposition 7.7 *For all $X, Y \subseteq V(G)$ with $Y \neq \varnothing$ and all $\alpha \in (1, 2]$, we have*

$$\sum_{m=0}^{\infty} \left(\frac{\alpha\Lambda}{\ln \alpha}\right)^{-m} bf^*_m(X, Y) \;\leq\; \alpha^{|Y|-1}\; . \tag{7.16}$$

Proof. Let $c = \alpha / \ln \alpha$. We shall show that

$$\sum_{m=0}^{M} (c\Lambda)^{-m}\, bf^*_m(X, Y) \;\leq\; \alpha^{|Y|-1} \tag{7.17}$$

for all $M \geq 0$, by using induction on $M + |X| + |Y|$. As noted above, we may assume that $X \cap Y = \varnothing$. The case $M = 0$ with X, Y arbitrary holds by the remark after the definition of $\mathcal{BF}^*_m(X, Y)$. The proposition also holds when $|X \cup Y| = 1$ and M is arbitrary, since $bf^*_m(\varnothing, \{y\}) = \delta_{m0}$. So assume $|X \cup Y| \geq 2$ and choose $y \in Y$. By Proposition 2.12, there exists a cocycle $C = E_G(L, R)$ in G and a vertex $z \in X \cup Y \setminus y$ such that $z \in L$, $X \cup Y \setminus z \subseteq R$, and $\sum_{e \in C} w_e \leq \Lambda$. Later we shall distinguish two cases, depending on whether z happens to lie in X or in Y.

For $H \in \mathcal{BF}^*_m(X, Y)$, let $F := F(H)$ be the set of edges of H that occur as the first edge in C on some path in H from z to $X \cup Y \setminus z$. Let H_1 be the connected



Figure 1: The graph $H \in \mathcal{BF}^*_m(X,Y)$. Blocks of H_1 and $H_2 = H - H_1$ are indicated schematically by large open circles; the edges of $C \cap E(H)$ are shown explicitly, with those in F drawn in bold. Vertices in X are indicated by small solid circles, vertices in Y are indicated by small open circles, and vertices in L^F, R^F by small solid squares.

component of $H \setminus F$ containing z, and let $H_2 = H \setminus V(H_1)$. Let L^F (resp. R^F) be the set of vertices of L (resp. R) that are incident with F; clearly $|L^F|, |R^F| \le |F|$. (This construction is illustrated in Figure 1.) For each nonempty $F \subseteq C$, let $\mathcal{BF}^*_m(X,Y;F)$ be the set of all subgraphs $H \in \mathcal{BF}^*_m(X,Y)$ such that $F(H) = F$.

Consider now the following two cases:

Case 1: $\boldsymbol{z \in X}$. Then $\mathcal{BF}^*_m(X,Y;\varnothing) = \varnothing$. We shall show that, for each nonempty $F \subseteq C$ with $|F| = f$ and each $H \in \mathcal{BF}_m(X,Y;F)$, we have $H_1 \in \mathcal{BF}^*_i(L^F,\{z\}) = \mathcal{BT}_i(L^F \cup \{z\})$ and $H_2 \in \mathcal{BF}^*_{m-f-i}(X \setminus z, Y \cup R^F)$ for some i $(0 \le i \le m-f)$. Note that by the above definitions $L^F \cup \{z\} \subseteq V(H_1)$ and $(X \setminus z) \cup (Y \cup R^F) \subseteq V(H_2)$.

We first consider H_1. Then H_1 is connected and $z \in V(H_1)$. Let B_1 be an end block of H_1. Then B_1 is either an end block of H, and hence $z \in \mathrm{Int}(B_1,H_1)$ or else B_1 is not an end block of H and hence $\mathrm{Int}(B_1,H_1) \cap L^F \ne \varnothing$. If H_1 is non-separable then either $V(H_1) = \{z\}$, or else H_1 contains z and at least one vertex of $L^F \setminus z$ (otherwise H_1 would be an end block of H with no internal vertex in $X \cup Y$). Thus $H_1 \in \mathcal{BF}^*_i(L^F,\{z\})$ for some i, $0 \le i \le m-f$.

We next consider H_2. Each component of H_2 is either a component of H, and hence contains a vertex of Y, or is not a component of H, and hence contains a vertex of R^F. Let B_2 be an end block of H_2. Then B_2 is either an end block of H, and hence satisfies $\mathrm{Int}(B_2, H_2) \cap [(X \setminus z) \cup Y] \neq \varnothing$, or is not an end block of H, and hence satisfies $\mathrm{Int}(B_2, H_2) \cap R^F \neq \varnothing$. Each isolated vertex of H_2 is either an isolated vertex of H, and hence belongs to Y, or is not an isolated vertex of H, and hence belongs to R^F. Let B_3 be an isolated block of H_2 which is not a single vertex. Then either: B_3 is an isolated block of H, and hence contains at least two vertices of $(X \setminus z) \cup Y$; or B_3 is an end block of H, and hence contains two distinct vertices, one in $(X \setminus z) \cup Y$ and the other in R^F; or B_3 is not a block of H, and hence contains two distinct vertices of R^F. Thus $H_2 \in \mathcal{BF}^*_{m-f-i}(X \setminus z, Y \cup R^F)$.

It follows that the map $H \mapsto (H_1, H_2)$ is a weight preserving injection from $\mathcal{BF}^*_m(X, Y; F)$ into $\bigcup_{i=0}^{m-f} \mathcal{BF}^*_i(L^F, \{z\}) \times \mathcal{BF}^*_{m-f-i}(X \setminus z, Y \cup R^F)$. Therefore,

$$bf^*_m(X,Y) \;\le\; \sum_{\varnothing \neq F \subseteq C} w(F) \sum_{i=0}^{m-f} bf^*_i(L^F, \{z\})\, bf^*_{m-f-i}(X \setminus z, Y \cup R^F)\;. \tag{7.18}$$

It follows that

$$\begin{aligned}
&\sum_{m=0}^{M} (c\Lambda)^{-m}\, bf^*_m(X,Y) \\
&\le\; \sum_{m=0}^{M} (c\Lambda)^{-m} \sum_{\varnothing \neq F \subseteq C} w(F) \sum_{i=0}^{m-f} bf^*_i(L^F, \{z\})\, bf^*_{m-f-i}(X \setminus z, Y \cup R^F) \\
&=\; \sum_{\varnothing \neq F \subseteq C} w(F)\, (c\Lambda)^{-f} \sum_{i=0}^{M-f} (c\Lambda)^{-i}\, bf^*_i(L^F, \{z\}) \sum_{j=0}^{M-f-i} (c\Lambda)^{-j}\, bf^*_j(X \setminus z, Y \cup R^F) \\
&\le\; \sum_{\varnothing \neq F \subseteq C} w(F)\, (c\Lambda)^{-f}\, \alpha^{|Y|-1+f} \qquad (7.19)
\end{aligned}$$

by the inductive hypothesis on $M + |X| + |Y|$. Using Lemma 6.2, we deduce that

$$\begin{aligned}
\sum_{m=0}^{M} (c\Lambda)^{-m}\, bf^*_m(X,Y) \;&\le\; \sum_{f=1}^{\infty} (c\Lambda)^{-f}\, \alpha^{|Y|-1+f} \sum_{\substack{F \subseteq C \\ |F| = f}} w(F) \\
&\le\; \sum_{f=1}^{\infty} (c\Lambda)^{-f}\, \alpha^{|Y|-1+f}\, \frac{\Lambda^f}{f!} \\
&=\; \alpha^{|Y|-1}(e^{\alpha/c} - 1) \;\le\; \alpha^{|Y|-1} \qquad (7.20)
\end{aligned}$$

since $c = \alpha / \ln \alpha$ and $\alpha \le 2$. Thus the proposition holds when $z \in X$.

Case 2: $\boldsymbol{z \in Y}$. Then $y, z \in Y$ and hence $|Y| \ge 2$. We can show in a similar way to Case 1 that, for each $F \subseteq C$ with $|F| = f$ and each $H \in \mathcal{BF}^*_m(X, Y; F)$, we have $H_1 \in \mathcal{BF}^*_i(L^F, \{z\})$ and $H_2 \in \mathcal{BF}^*_{m-f-i}(X, (Y \setminus z) \cup R^F)$ for some i $(0 \le i \le m - f)$. [Let us remark that if $F = \varnothing$, then H_1 is a component of H consisting of just the isolated vertex z.] The map $H \mapsto (H_1, H_2)$ is an injection from $\mathcal{BF}^*_m(X, Y; F)$ into $\bigcup_{i=0}^{m-f} \mathcal{BF}^*_i(L^F, \{z\}) \times \mathcal{BF}^*_{m-f-i}(X, (Y \setminus z) \cup R^F)$. Thus

$$bf^*_m(X, Y) \le \sum_{F \subseteq C} w(F) \sum_{i=0}^{m-f} bf^*_i(L^F, \{z\}) \, bf^*_{m-f-i}(X, (Y \setminus z) \cup R^F) \,. \tag{7.21}$$

Hence

$$\begin{aligned}
&\sum_{m=0}^{M} (c\Lambda)^{-m} \, bf^*_m(X, Y) \\
&\le \sum_{m=0}^{M} (c\Lambda)^{-m} \sum_{F \subseteq C} w(F) \sum_{i=0}^{m-f} bf^*_i(L^F, \{z\}) \, bf^*_{m-f-i}(X, (Y \setminus z) \cup R^F) \\
&= \sum_{F \subseteq C} w(F) \, (c\Lambda)^{-f} \sum_{i=0}^{M-f} (c\Lambda)^{-i} \, bf^*_i(L^F, \{z\}) \sum_{j=0}^{M-f-i} (c\Lambda)^{-j} \, bf^*_j(X, (Y \setminus z) \cup R^F) \\
&\le \sum_{F \subseteq C} w(F) \, (c\Lambda)^{-f} \, \alpha^{|Y|-2+f} \qquad (7.22)
\end{aligned}$$

by the inductive hypothesis on $M + |X| + |Y|$. (Note that when $F = \varnothing$, we have $L^F = R^F = \varnothing$ and hence $|(Y \setminus z) \cup R^F| = |Y| - 1$.) Using Lemma 6.2, we deduce that

$$\begin{aligned}
\sum_{m=0}^{M} (c\Lambda)^{-m} \, bf^*_m(X, Y) &\le \sum_{f=0}^{\infty} (c\Lambda)^{-f} \, \alpha^{|Y|-2+f} \sum_{\substack{F \subseteq C \\ |F| = f}} w(F) \\
&\le \sum_{f=0}^{\infty} (c\Lambda)^{-f} \, \alpha^{|Y|-2+f} \, \frac{\Lambda^f}{f!} \\
&= \alpha^{|Y|-2} e^{\alpha/c} \;=\; \alpha^{|Y|-1} \qquad (7.23)
\end{aligned}$$

since $c = \alpha / \ln \alpha$. Thus the proposition holds when $z \in Y$. $\square$

Remark. The proof technique of Case 1 can be used to prove the induction hypothesis

$$\sum_{m=0}^{M} (c\Lambda)^{-m} bf^*_m(X,Y) \;\le\; \alpha^{|Y|-1} \tag{7.24}$$

whenever $e^{\alpha/c} \le 2$. (In particular, it can handle the apparently best-possible values $c = 1/\ln 2$ and $\alpha = 1$.) Likewise, the proof technique of Case 2 can be used to prove the induction hypothesis (7.24) whenever $e^{\alpha/c} \le \alpha$. The trouble is that we need the same hypothesis to work for both cases, since we don't know *a priori* whether z will lie in X or in Y. Therefore, the best we can do — at least with this proof technique — seems to be to choose $\alpha \in (1,2]$ and then set $c = \alpha/\ln\alpha$.

Example 7.4. Let T be any tree (e.g. a long path would do), and let G be the graph $T^{(s)}$ obtained from T by replacing each edge by s parallel edges. Let all edge weights w_e equal Λ/s, so that the maxmaxflow is Λ. Let $x, y \in V(T^{(s)})$ with $\operatorname{dist}(x,y) = \ell$. Then $bf_m(\{x\},\{y\}) = bf^*_m(\{x\},\{y\})$, and we have

$$\sum_{m=0}^{\infty} \zeta^m bf_m(\{x\},\{y\}) \;=\; \left[\left(1 + \frac{\Lambda}{s}\zeta\right)^{s} - 1 \right]^{\ell} \;\overset{s\to\infty}{\longrightarrow}\; (e^{\Lambda\zeta} - 1)^{\ell} \,. \tag{7.25}$$

Since ℓ can be arbitrarily large, a universal upper bound on $\sum_{m=0}^{\infty} \zeta^m bf_m(X,Y)$ is impossible for $\zeta > (\ln 2)/\Lambda$, even when $|X| = |Y| = 1$. $\square$

Example 7.5. Let G be the union of k disjoint copies of $K^{(s)}_{1,r}$, with all edge weights $w_e = \Lambda/s$. Let X be the central vertices and Y the remaining vertices, so that $|X| = k$ and $|Y| = kr$. Then

$$\sum_{m=0}^{\infty} \zeta^m bf_m(X,Y) \;=\; r^k \left[\left(1 + \frac{\Lambda}{s}\zeta\right)^{s} - 1 \right]^{k} \;\overset{s\to\infty}{\longrightarrow}\; r^k (e^{\Lambda\zeta} - 1)^{k} \tag{7.26}$$

and

$$\sum_{m=0}^{\infty} \zeta^m bf^*_m(X,Y) \;=\; \left[\left(1 + \frac{\Lambda}{s}\zeta\right)^{rs} - 1 \right]^{k} \;\overset{s\to\infty}{\longrightarrow}\; (e^{r\Lambda\zeta} - 1)^{k} \,. \tag{7.27}$$

Thus, if any universal upper bound is possible for $\zeta = (\ln 2)/\Lambda$, the right-hand side has to be at least $(|Y|/|X|)^{|X|}$ for bf and $(2^{|Y|/|X|} - 1)^{|X|}$ for bf^*. $\square$

These examples suggest the following conjectures:

Conjecture 7.8 *For all $X, Y \subseteq V(G)$ with $Y \neq \varnothing$, we have*

$$\sum_{m=0}^{\infty} \left(\frac{\Lambda}{\ln 2}\right)^{-m} bf_m(X,Y) \;\le\; |Y|^{|X|} \,. \tag{7.28}$$

Conjecture 7.9 *For all $X, Y \subseteq V(G)$ with $Y \neq \varnothing$, we have*

$$\sum_{m=0}^{\infty} \left(\frac{\Lambda}{\ln 2} \right)^{-m} bf_m^*(X,Y) \;\le\; 2^{|Y|} - 1 \;. \tag{7.29}$$

Conjecture 7.10 (a special case of Conjecture 7.8 or 7.9) *For all nonempty $X \subseteq V(G)$, we have*

$$\sum_{m=0}^{\infty} \left(\frac{\Lambda}{\ln 2} \right)^{-m} bt_m(X) \;\le\; 1 \;. \tag{7.30}$$

7.2 The class $\mathcal{B}_m(X)$

We conclude by discussing a larger class of block forests, which is roughly the block analogue of the class $\mathcal{H}_m(X)$ considered in Section 5.2. For $X \subseteq V(G)$, let $\mathcal{B}_m(X)$ be the set of all m-edge subgraphs H in G (connected or not) such that

(B1) $X \subseteq V(H)$;

(B2) each end block of H contains at least one element of X as an internal vertex [that is, $\mathrm{Int}(B,H) \cap X \neq \varnothing$]; and

(B3) each isolated block of H is either an isolated vertex belonging to X or else contains at least two vertices of X.

[It follows from (B2) and (B3) that each component of H is either an isolated vertex belonging to X or else contains at least two vertices of X. In particular, $\mathcal{B}_m(X) \subseteq \mathcal{C}_m(X)$. Note also that $\mathcal{BF}_m(X,Y) \subseteq \mathcal{BF}_m^*(X,Y) \subseteq \mathcal{B}_m(X \cup Y)$.] Note the following special cases:

- For any X, $\mathcal{B}_0(X)$ has as its single element the edgeless graph with vertex set X.
- $\mathcal{B}_m(\varnothing) = \varnothing$ for $m \ge 1$.
- $\mathcal{B}_m(\{x\}) = \varnothing$ for $m \ge 1$.
- $\mathcal{B}_m(\{x,y\})$ is the set of m-edge xy-block paths when $x \neq y$ and $m \ge 1$. [Hence $\mathcal{B}_m(\{x,y\}) = \mathcal{BT}_m(\{x,y\}) = \mathcal{BF}_m(\{x\},\{y\}) = \mathcal{BF}_m^*(\{x\},\{y\})$.]

Recall that $w(H) = \prod_{e \in E(H)} w_e$. Define the weighted counts

$$b_m(X) \;=\; \sum_{H \in \mathcal{B}_m(X)} w(H) \;. \tag{7.31}$$

We then have the following bound in terms of maxmaxflow:

Proposition 7.11 *Whenever* $|X| = k \geq 1$ *we have*

$$b_m(X) \;\leq\; B(m,k)\,\Lambda^m \tag{7.32}$$

where

$$B(m,k) \;=\; 2^m C(m, \frac{k-1}{2}) \;=\; \begin{cases} (k-1)(2m+k-1)^{m-1}/m! & \textit{for } k \neq 1 \\ \delta_{m0} & \textit{for } k = 1 \end{cases} \tag{7.33}$$

For the case of greatest interest, namely $\mathcal{B}_m(\{x,y\}) = \mathcal{BF}^*_m(\{x\},\{y\})$, the bound of Proposition 7.11 behaves roughly like $(2e\Lambda)^m$, which is much worse than the bound $(2\Lambda/\ln 2)^m$ of Propositions 7.2 and 7.7. However, as we shall see, the proof of Proposition 7.11 is quite a bit simpler than that of Proposition 7.7.

Before beginning the proof of Proposition 7.11, let us note some facts about the numbers $B(m,k)$, which follow easily from the facts about the $C(m,k)$ already discussed in Section 6:

(a) For each integer $m \geq 0$, $B(m,k)$ is an increasing function of $k \geq 1$.

(b) Generating function: If $\mathsf{C}(z)$ solves the equation (6.4), then

$$\mathsf{C}(2z)^{(k-1)/2} \;=\; \sum_{m=0}^{\infty} B(m,k)\, z^m \tag{7.34}$$

This is an immediate translation of (6.5).

(c) For all k_1, k_2, m we have

$$\sum_{i=0}^{m} B(i,k_1)\, B(m-i,k_2) \;=\; B(m, k_1+k_2-1)\;. \tag{7.35}$$

This is an immediate consequence of (7.34).

(d) For all k,

$$B(m,k) \;=\; \sum_{f=0}^{m} \frac{1}{f!}\, B(m-f, k-1+2f)\;. \tag{7.36}$$

This is an immediate translation of the identity (6.7) with $z = 1/2$.

(e) For each fixed $k > 1$, we have

$$B(m,k) \;=\; (2e)^m m^{-3/2} \frac{(k-1)e^{(k-1)/2}}{\sqrt{8\pi}}\,[1 + O(1/m)] \tag{7.37}$$

as $m \to \infty$. This is an immediate translation of (6.9).

PROOF OF PROPOSITION 7.11. The Proposition holds trivially when $k = 1$, so we may assume that $k \geq 2$. Choose $x_1, x_2 \in X$ and let C be a minimum cut in G separating x_1 from x_2, so that $\sum_{e \in C} w_e \leq \Lambda$. Let G_1 be the component of $G - C$ containing x_1, and let G_2 be the union of the remaining components of $G - C$. By construction, we have $x_1 \in V(G_1)$ and $x_2 \in V(G_2)$, and each edge in C joins a vertex of G_1 to a vertex of G_2. Let $X_i = X \cap V(G_i)$ and $k_i = |X_i|$ for $i = 1, 2$. Since $x_1 \in X_1$ and $x_2 \in X_2$, we have $1 \leq k_i \leq k - 1$ for $i = 1, 2$; and of course $k_1 + k_2 = k$.

We shall classify the subgraphs H of G belonging to $\mathcal{B}_m(X)$ according to $F \equiv E(H) \cap C$. So, for each $F \subseteq C$, let us set $f = |F|$; define Y_i^F $(i = 1, 2)$ to be the set of vertices of G_i that are incident with F (note that $|Y_i^F| \leq f$); and finally, let $\mathcal{B}_m(X; F)$ be the set of all $H \in \mathcal{B}_m(X)$ that have $E(H) \cap C = F$.

For $H \in \mathcal{B}_m(X; F)$, we let $H_i = H \cap G_i$ and $m_i = |E(H_i)|$; note that $H = H_1 \cup H_2 \cup F$ and $m = m_1 + m_2 + f$. We shall show that $H_i \in \mathcal{B}_{m_i}(X_i \cup Y_i^F)$ for $i = 1, 2$. (The argument is illustrated in Figure 2.) Note first that since $X \subseteq V(H)$ we have $X_i \subseteq V(H_i)$. Furthermore, each end block B of H_i is either an end block of H or else satisfies $\mathrm{Int}(B, H_i) \cap Y_i^F \neq \varnothing$. Hence $\mathrm{Int}(B, H_i) \cap (X_i \cup Y_i^F) \neq \varnothing$. Each isolated vertex v of H_i is either an isolated vertex of H or else belongs to Y_i^F. Thus $v \in X_i \cup Y_i^F$. Finally, each isolated block B of H_i that is not an isolated vertex of H_i is either an isolated block of H, or else an end block of H with its cut vertex in Y_i^F, or else satisfies $|V(B) \cap Y_i^F| \geq 2$. Thus $|V(B) \cap (X_i \cup Y_i^F)| \geq 2$.

It follows that we can construct a weight-preserving [except for a factor $w(F)$] bijection of $\mathcal{B}_m(X; F)$ onto a subset of $\bigcup_{i=0}^{m-f} \mathcal{B}_i(X_1 \cup Y_1^F) \times \mathcal{B}_{m-f-i}(X_2 \cup Y_2^F)$. Thus

$$b_m(X) \;\leq\; \sum_{F \subseteq C} w(F) \sum_{i=0}^{m-f} b_i(X_1 \cup Y_1^F)\, b_{m-f-i}(X_2 \cup Y_2^F)\;. \tag{7.38}$$

Since $i \leq m - f$, $|Y_1^F| \leq f$ and $|X_1| < k$, we have $i + |X_1 \cup Y_1^F| < m + k$; and likewise we have $(m - f - i) + |X_2 \cup Y_2^F| < m + k$. Therefore, we can use induction on $m + k$ (the Proposition being true for the initial case $m + k = 1$) to deduce that

$$\begin{aligned}
b_m(X) \;&\leq\; \sum_{F \subseteq C} w(F) \sum_{i=0}^{m-f} \Lambda^{m-f}\, B(i, |X_1 \cup Y_1^F|)\, B(m - f - i, |X_2 \cup Y_2^F|) \\
&\leq\; \sum_{F \subseteq C} w(F) \sum_{i=0}^{m-f} \Lambda^{m-f}\, B(i, k_1 + f)\, B(m - f - i, k_2 + f) \\
&\leq\; \sum_{f=0}^{m} \frac{\Lambda^f}{f!} \sum_{i=0}^{m-f} \Lambda^{m-f}\, B(i, k_1 + f)\, B(m - f - i, k_2 + f) \\
&=\; \Lambda^m B(m, k)\;,
\end{aligned} \tag{7.39}$$



Figure 2: The graph $H \in \mathcal{B}_m(X)$. Blocks of H_1 and H_2 are indicated schematically by large open circles; the edges of F are shown explicitly. Vertices in X are indicated by small solid circles, and vertices in Y_1^F, Y_2^F are indicated by small solid squares.

where the second line used the fact that $B(m,k)$ increases with k for $k \geq 1$, the third line used Lemma 6.2, and the last line used identities (7.35) and (7.36) and the fact that $k_1 + k_2 = k$. $\square$

Corollary 7.12 *Fix a weighted graph* $(G, \mathbf{w})$ *and an edge* $e \in E(G)$*. Then the sum of the weights of the 2-connected* m*-edge subgraphs of* G *containing* e *is at most* $B(m-1,2)\, \Lambda(G-e,\mathbf{w})^m$*.*

PROOF. If $e = xy$, this follows immediately from Proposition 7.11 applied to $G - e$ by taking $X = \{x, y\}$. $\square$

8 A Conjecture on Chromatic Roots

As mentioned in the Introduction, our principal motivation for studying maxmaxflow is the conjecture that it can be used to bound the roots of chromatic polynomials $P_G(q)$:

Conjecture 8.1 *There exist universal constants $C(\Lambda) < \infty$ such that all the chromatic roots (real or complex) of all loopless graphs of maxmaxflow Λ lie in the disc $|q| \le C(\Lambda)$. Indeed, we conjecture that $C(\Lambda)$ can be taken to be linear in Λ.*

The analogous theorem using maximum degree was proven in [12]: the approach taken there is to decompose a spanning subgraph of G into its connected components and to treat these components as a "polymer gas". The desired bound on chromatic roots then follows from standard bounds on the zeros of a polymer-gas partition function, once one has an exponential bound in terms of maximum degree on the number of connected m-edge subgraphs of G containing a specified vertex (Proposition 6.1 of the present paper or the variants thereof contained in [12, Section 4.2]). Moreover, by special arguments it was possible to bound the chromatic roots also in terms of second-largest degree [12, Section 6]. We are now trying to prove Conjecture 8.1 by modifying the arguments of [12] to decompose a spanning subgraph of G into its *2-connected* components. Although the pointwise bound implied by Corollary 7.4 of the present paper has a presumably non-optimal constant — unlike Proposition 6.1 and its relatives, which are sharp — it should nevertheless be good enough to prove Conjecture 8.1 provided that the other difficulties (such as controlling the interaction between the 2-connected components) can be overcome.

Acknowledgments

We wish to thank Jan van den Heuvel and Akira Saito for valuable conversations. We also thank Gordon Royle for drawing our attention to references [1, 8], and for many valuable conversations.

This research was supported in part by U.S. National Science Foundation grants PHY–9900769, PHY–0099393 and PHY–0424082. It was begun while one of the authors (A.D.S.) was a Visiting Fellow at All Souls College, Oxford, where he was hosted by the Department of Theoretical Physics and supported in part by U.K. Engineering and Physical Sciences Research Council grant GR/M 71626.

References

[1] Bollobás, B. (1966) On graphs with at most three independent paths connecting any two vertices. *Studia Sci. Math. Hungar.* **1**, 137–140.

[2] Bollobás, B. (1978) *Extremal Graph Theory*. Academic Press, London.

[3] Bollobás, B. (1998) *Modern Graph Theory*. Springer-Verlag, New York.

[4] Gomory, R.E. and Hu, T.C. (1961) Multi-terminal network flows. *J. Soc. Indust. Appl. Math.* **9**, 551–570. Reprinted in D.R. Fulkerson (editor), *Studies in*

Graph Theory, Part I, Studies in Mathematics #11, pp. 172–199 (Mathematical Association of America, Washington, 1975).

[5] Leonard, J.L. (1972) On graphs with at most four line-disjoint paths connecting any two vertices. *J. Combin. Theory B* **13**, 242–250.

[6] Leonard, J.L. (1973) Graphs with 6-ways. *Canad. J. Math.* **25**, 687–692.

[7] Lovász, L. and Plummer, M.D. (1986) *Matching Theory*. North-Holland Mathematics Studies #121. Annals of Discrete Mathematics #29. North-Holland, Amsterdam-New York / Akadémiai Kiadó, Budapest.

[8] Mader, W. (1973) Ein Extremalproblem des Zusammenhangs von Graphen. *Math. Z.* **131**, 223–231.

[9] Mader, W. (1979) Connectivity and edge-connectivity in finite graphs. In B. Bollobás (editor), *Surveys in Combinatorics* (Proceedings of the Seventh British Combinatorial Conference), London Mathematical Society Lecture Note Series #38, pp. 66–95. Cambridge University Press, Cambridge.

[10] Shrock, R. and Tsai, S.-H. (1998) Ground state degeneracy of Potts antiferromagnets: Cases with noncompact W boundaries having multiple points at $1/q = 0$. *J. Phys. A* **31**, 9641–9665, cond-mat/9810057 at arXiv.org.

[11] Shrock, R. and Tsai, S.-H. (1999) Ground state degeneracy of Potts antiferromagnets: Homeomorphic classes with noncompact W boundaries. *Physica A* **265**, 186–223, cond-mat/9811410 at arXiv.org.

[12] Sokal, A.D. (2001) Bounds on the complex zeros of (di)chromatic polynomials and Potts-model partition functions. *Combin. Probab. Comput.* **10**, 41–77, cond-mat/9904146 at arXiv.org.

[13] Sokal, A.D. (2004) Chromatic roots are dense in the whole complex plane. *Combin. Probab. Comput.* **13**, 221–261, cond-mat/0012369 at arXiv.org.

[14] Thomassen, C. (1997) The zero-free intervals for chromatic polynomials of graphs. *Combin. Probab. Comput.* **6**, 497–506.